Pepper Ranch Preserve

Land Management Plan (5 Yr. Revision)



Formatted: Font: 24 pt
Formatted: Font: 24 pt
Formatted: Font: 24 pt

Managed by: Conservation Collier Program
Collier County, Florida
May 2011 – May 2021 (10-yr plan)
Prepared by:

Johnson Engineering & Conservation Collier Staff;

Collier County Parks and Recreation Facilities Management Division epartment



Formatted: Normal, Space Before: 6 pt, After: 6 pt

G:\Conservation Collier\Land Management\PepperRanchPreserve\Land Management Plans\FinalManagementPlan\PRP_5 vear revisionFinal_EMP_April11.do

Pepper Ranch Preserve

Land Management Plan Executive Summary

Lead Agency: Conservation Collier Program

Properties included in this Plan: Pepper Ranch Preserve

Preserve lands consist of ten contiguous parcels located within Sections 22, 26, 27, 28, 33 & 35, Township 46 South, Range 28 East, in Collier County. A property survey and full legal description is provided in Appendix 1.

Parcel Foli	o Numbers
00052360002	00052680009
00052440003	00054040003
00053000002	00053920001
00052960004	00053840000
00052640007	00053200006

Original Acreage Breakdown:

Natural Community	<u>Acreage</u>
Improved pasture	619.64
Depression marsh, disturbed	311.29
Upland mixed forest	270.92
Slough	243.38
Bottomland forest	241.63
Upland mixed forest, disturbed	160.59
Mesic flatwoods	149.95
Strand swamp	82.41
Mesic flatwoods, disturbed	77.20
Improved pasture, hydric	47.70
Dry prairie, disturbed	45.09
Wet flatwoods	43.66
Depression marsh	42.60
Ditches	35.15
Upland mixed forest, burned	34.65
Shell road, graded and drained	20.63
Slough, disturbed	14.17
Prairie hammock	11.22

Formatted: Font: 6 pt

Formatted: Normal, Space Before: 6 pt, After: 6 pt **Formatted:** Left, Indent: Left: 1.5", First line: 0.5"

Formatted: Justified, Indent: Left: 0.25", First line: 0"

Commented [S1]: Will these be combined?

Spoil, exotics	9.49
Natural Community (continued)	Acreage
Wet prairie, disturbed	7.99
Bottomland forest, disturbed	6.27
Prairie hammock, disturbed	5.88
Oil field	5.63
Wet prairie	5.21
Primitive trail	4.90
Brazilian pepper, hydric	4.26
Dry prairie	3.53
Borrow pond	1.43
Spoil	1.24
Australian pine	1.15
Brazilian pepper	1.15
TOTAL	2510.01

Management Responsibilities:

Agency: Collier County - Conservation Collier Program

Preserve Manager: Designated Collier County Environmental Specialist

Designated Land Use: Preservation

Unique Features: Largest Conservation Collier acquisition to date with frontage on north side of Lake Trafford

Archaeological/Historical: The Pepper Ranch Preserve is within an area of historical and archaeological probability. One prehistoric and archaeological site has already been discovered near Lake Trafford and the <u>Ledge/visitor</u> center is considered a historic structure. Other historical and archaeological sites are most likely present on the property. The County will notify the Division of Historical Resources immediately if evidence is discovered to suggest that any additional cultural resources are present.

Management Needs:

Monitoring of biological resources;

Exotic plant removal and maintenance

Conduct a hydrological analysis of the preserve to better determine restoration needs;

Possoble Possible restoration of select areas after exotic removal;

Implementation of a prescribed fire management program;

Habitat management to enhance protection of native and listed species populations; and

Monitoring Planning for public use.

Public Involvement: Workeding with Immokalee Civic Group(s) including the Immokalee Community Redevelopment Association, Collier County Sheriff's Department and neighboring property owners to discuss public use and access issues. Two public meetings were held (January 28 and February 11, 2010) to provide the general public an opportunity to review and comment on this Pepper Ranch Preserve Land Management Plan. Specifically, these meetings gave the general public an opportunity to learn about the Conservation Collier Program, future land management plans and potential for public use at Pepper Ranch Preserve. This forum also allowed the public to voice any concerns or objections they may have had with any of these issues as presented here in the Final Land Management Plan.

Conservation Collier and the Collier County Board of County Commissioners (BCC) held workshops (February 18 and March 18, 2010) with outdoor sportsmen's clubs and hunters to develop a Hunt Program for Pepper Ranch Preserve. Over 20 people attended the February 18th, 2010 meeting, including representatives from the Florida Fish and Wildlife Conservation Commission (FWC) and Commissioner Colletta, to discuss how best to create a Hunt Program for the preserve. A significant amount of information was collected from this meeting to incorporate into a Hunt Management Plan for the preserve including how often to allow hunts, how many hunters, what types of weapons, check station and registration logistics etc.

After the February 18th, 2010 meeting, the first draft of the Hunt Management Plan was created and posted for public review. During the March 18th, 2010 meeting, approximately 10 local hunters attended including Commissioner Colletta, to review the draft Hunt Management Plan. The consensus at this meeting was to divide the ranch into sections and each hunter would have to stay in their section during the hunt. The use of rifles was also discussed as well as safety zones and parking areas.

Over the weekend of April 16-18, 2010, the first Annual Youth Hunt was held at Pepper Ranch Preserve. FWC and volunteers ran the hunt and it was a huge success. There were 15 kids that participated age 12-17 and 4 hogs were harvested.

At the April 27, 2010 Collier County BCC meeting, the draft Public Hunt Management Plan for Pepper Ranch was brought for BCC approval. At this meeting, Conservation Collier staff was directed to hold two (2) additional public meetings to determine if hunting should be limited to youth hunts only or to the general public. These meetings were held on May 12 and May 19, 2010. The results of that meeting were to hold at least 2 youth hunts per year and -leave the remaining hunts open to the general public.

Two additional public meetings were held to provide the public an opportunity to give input on the required 5-year plan update. The meetings were held on March 21, 2017 in Immokalee, and the second was held at the Collier County Extension office on March 28, 2017.

Management Goals:

- Goal 1: Eliminate or significantly reduce human impacts to indigenous flora and fauna
- Goal 2: Develop a baseline monitoring program
- **Goal 3:** Remove or control populations of invasive, exotic or problematic flora and fauna to restore and maintain natural habitats
- Goal 4: Create a Prescribed Fire Management Plan
- Goal 5: Restore native vegetation as needed
- Goal 6: Monitor Develop a plan for public use
- Goal 7: Facilitate uses of the site for educational purposes
- Goal 8: Provide a plan for security and disaster preparedness
- **Goal 9:** Provide preliminary panther habitat unit (PHU) calculations and a draft monitoring plan per U.S. Fish and Wildlife Service (USFWS) requirements for an onsite panther conservation bank

TABLE OF CONTENTS

		TABLE OF CONTENTS	Page
			Tage
1.0	Intro	duction	9
	1.1	Conservation Collier: Land Acquisition Program and Management	
		Authority	10
	1.2	Purpose and Scope of the Land Management Plan	
	1.3	Location of the Pepper Ranch Preserve	10
	1.4	Regional Significance of the Pepper Ranch Preserve	1
	1.5	Nearby Public Lands and Designated Water Resources	
	1.6	Public Involvement	
2.0	Natu	ral Resources	14
	2.1	Physiography	
		2.1.1 Topography and Geomorphology	
		2.1.2 Geology	
		2.1.3 Soils	
		2.1.4 Hydrology/Water Management	
	2.2	Climate	
	2.3	Natural Plant Communities	
		2.3.1 Uplands: Upland Mixed Forest	
		2.3.2 Wetlands: Depression Marsh	
		2.3.3 Wetlands: Slough	
		2.3.4 Wetlands: Bottomland Forest	
		2.3.5 Uplands: Mesic Flatwoods	
		2.3.6 Other Natural Communities	30
		2.3.7 Altered Communities	
	2.4	Native Plant and Animal Species	33
		2.4.1 Plant Species	
		2.4.2 Animal Species	
	2.5	Listed Species	3
		2.5.1 Listed Plant Species	
		2.5.2 Listed Animal Species	
	2.6	Invasive, Non-native and Problem Species	
		2.6.1 Invasive and Problem Plant Species	
		2.6.2 Invasive and Problem Animal Species	
3.0	Previ	ious Conditions of the Preserve; Current Use of the Preserve and	
0.0		cent Land Uses	4
	3.1	Previous Land Uses of the Preserve	
	3.2	Previous Land Uses of Adjoining Properties	
	3.3	Current Land Uses of the Preserve	
	3.4	Current Land Uses of Adjoining Properties	
	3.5	Cultural, Historical and Archeological Resource Protection	
	3.6	Major Accomplishments since Acquisition	

4.0 F	uture Use of the Pepper Ranch Preserve including Management Issues,	
G	oals and Objectives	
4.	.1 Management Plan Framework	54
	4.1.1 Preserve Manager: Contact Information	
4.	.2 Planned Uses and Assessment of their Impacts	54
	4.2.1 Identification of Public Uses Consistent with Preservation,	
	Enhancement, Restoration, Conservation and Maintenance of	
	the Resources	54
4.	.3 Desired Future Conditions	56
4.	.4 Goals for the 10 year period 2010-2020	57
4.	.5 Establish an Operational Plan for the Pepper Ranch Preserve	
	4.5.1 Maintenance	
	4.5.2 Estimated Annual Costs and Funding Sources	83
	4.5.3 Potential for Contracting Restoration and Management	
	Activities by Private Vendors	87
5.0 L	iterature Cited	
	<u>TABLES</u>	
Table 1	Acquisition History and Status of Pepper Ranch Preserve	
Table 2	Public Lands Located near the Pepper Ranch Preserve	13
Table 3	Extent of Florida Land Use, Cover and Forms Classification System	
	Designations from 2009 on the Pepper Ranch Preserve	20
Table 4	Summary of Natural Communities on the Pepper Ranch Preserve	
Table 5	Faunal Species Observed at Pepper Ranch Preserve	34
Table 6	Breeding Bird Species Recorded in the Corkscrew and Immokalee	
	Quadrangles Encompassing the Pepper Ranch Preserve	
Table 7	Listed Plant Species Detected at the Pepper Ranch Preserve	
Table 8	Non-Indigenous and Invasive Plant Species at Pepper Ranch Preserve	
Table 9	Major Accomplishments during previous years	
Table 10	, I	
	FLEPPC Category I species	
	Panther Habitat Unit Calculations including SSA 7 Pre Restoration	
	<u>11</u>	
	Panther Habitat Unit Calculations including SSA Post Restoration	
	Panther Habitat Unit Calculations excluding SSA - Post Restoration	
	<u>12</u>	
Table 16	<u>13</u>	Estimated Annual Land Management Budget
Table 17	14	Potential Contracting for Restoration and Manageme

FIGURES

Figure 1	General Location of and Directions to Pepper Ranch Preserve	11
Figure 2	Conserved Lands in Collier County, Florida Including Lands Owned by	
U	Conservation Collier.	12
Figure 3	Preserves and Protected Lands in the Vicinity of Pepper Ranch Preserve	
Figure 4	Aerial View of the Pepper Ranch Preserve	
Figure 5	Soil Units on the Pepper Ranch Preserve	
Figure 6	Distribution of Natural Communities and other Land Uses on the Pepper	
C	Ranch Preserve; 2009 FLUCFCS Layer	22
Figure 7	Extent of Natural Plant Communities Currently Found on the Pepper	
C	Ranch Preserve.	24
Figure 8	1940 Aerial View of Pepper Ranch Preserve	
Figure 9	1953 Aerial View of Pepper Ranch Preserve	
Figure 10	1963 Aerial View of Pepper Ranch Preserve	
Figure 11	Pepper Ranch Easement and Overlay Map	
Figure 12	Conceptual Site Plan for Pepper Ranch Preserve	
	•	
	<u>APPENDICIES</u>	
Appendix 1	Pepper Ranch Preserve Legal Description	
Appendix 2	Florida Land Use, Cover and Forms Classification System Designations	
	for Pepper Ranch Preserve – (2'x3' map)	
Appendix 3	Florida Natural Areas Inventory Designations for Pepper Ranch Preserve – (2'x3' map)	
Appendix 4	Floristic Inventory of Pepper Ranch Preserve	
Appendix 5	Management Unit Map of Pepper Ranch Preserve	
Appendix 6	Cattle Lease	
Appendix 7	Pepper Ranch Preserve – Quality Wildlife Management Hunt Program	
Appendix /	2010 and 2011 Brochure	
Appendix 8	Pepper Ranch Preserve – Land Use Compatibility Matrix	
* *		
Appendix 9	Parcel Folio Map of Pepper Ranch Preserve	

1.0 Introduction

The Pepper Ranch Preserve is 2,510.01 acres of natural and agricultural lands located along the north shore of Lake Trafford in north central Collier County, Florida. The preserve contains various native plant communities, including bottomland forest, upland mixed forest, strand swamp, slough, pine flatwoods, depression marshes, and improved pastures.

A site assessment to determine compliance with the Conservation Collier initial screening criteria was conducted in September, October and November 2007 and the Conservation Collier Program purchased the property in February 2009. The County holds fee simple title to the Pepper Ranch Preserve. The Conservation Collier program manages these lands under authority granted by the Conservation Collier Ordinance 2002-63 as amended (2007-65) (available from www.municode.com). Initial acquisition activities are summarized in Table 1.

	Table 1: Acquisition History and Status of Pepper Ranch Preserve		
Year	Benchmark		
2006	FLUCFCS mapping of a portion of the preserve conducted by Scheda		
2007	Property nominated to the Conservation Collier Program		
2007	Initial Site Assessment by Conservation Collier Staff		
2007	Acceptance of Initial Criteria Screening Report by the Conservation Collier Land Acquisition Advisory Committee		
2008	Phase I Environmental Assessment Conducted by Environmental Consulting and Technology, Inc. for Collier County		
2008	Approved for purchase by the Board of County Commissioners (BCC)		
2009	Purchase of the Pepper Ranch property		
2009	Developed Interim Management Plan		
2009	BCC approved the Interim Management Plan		
2010	Completed Final Land Management Plan		
2011	BCC approved Final Management Plan		
<u>2017</u>	Completed Final Management Plan 5 year update		

The preserve consists of approximately 43% (± 1087.15 acres) wetland plant communities and approximately 57% (± 1422.86 acres) upland plant communities. Conservation, restoration and natural resource-based recreation are the designated uses of this property. Management activities allowed include those necessary to preserve, restore, secure and maintain this environmentally sensitive land for the benefit of present and future generations. Public use of the site must be consistent with these management goals.

This is the Final Management Plan for the Pepper Ranch Preserve. This 10-year management plan was approved by ill be submitted to the Collier County Board of County Commissioners (BCC) for its approval on 6/28/11. Twhen approved, this plan

will-replaced the Interim Management Plan. Changes made to this plan during the 5-year review process will be brought before the BCC for their approval in-2017.

1.1 Conservation Collier: Land Acquisition Program and Management Authority

The Conservation Collier program was originally approved by voters in November 2002 and subsequently confirmed in the November 2006 ballot referendum. Both voter-approved referendums enable the program to acquire environmentally sensitive conservation lands within Collier County, Florida (Ordinance 2002-63, as amended 2007-65). Properties must support at least two of the following qualities to qualify for further consideration: rare habitat, aquifer recharge, flood control, water quality protection, and listed species habitat. The BCC appointed a Land Acquisition Advisory Committee to consider any selected or nominated properties that an owner has indicated a willingness to sell. The committee recommends property purchases for final approval by the BCC.

Lands acquired with Conservation Collier funds are titled to "COLLIER COUNTY, a political subdivision of the State of Florida, by and through its Conservation Collier program." The Collier County BCC established the Conservation Collier program to implement the program and to manage acquired lands. As such, Conservation Collier holds management authority for the Pepper Ranch Preserve. Conservation Collier is therefore established to acquire, preserve, restore, and maintain vital and significant threatened natural lands, forest, upland and wetland communities located in Collier County, for the benefit of present and future generations.

1.2 Purpose and Scope of the Land Management Plan

The purpose of the plan is to provide management direction for Pepper Ranch Preserve by identifying the goals and objectives necessary to eliminate or minimize any threats to the resources and integrity of the preserve. This text is a working document that establishes the foundation of the ten-year plan by identifying the appropriate management techniques necessary to preserve and/or restore the resource.

This plan will balance resource restoration and protection with natural resource-based recreational and educational use while looking at restoration needs, listed species protection and maintenance of the site free of invasive, exotic plant and animal species. This plan is divided into sections that incorporate an introduction, descriptions of the natural and cultural resources, projected uses of the property, management issues, and goals and objectives.

1.3 Location of the Pepper Ranch Preserve

Pepper Ranch Preserve is located at 6315 Pepper Road just west of the town of Immokalee, Florida (See Figure 1; legal description in Appendix 1). It is in north central Collier County, along the north shoreline of Lake Trafford within Sections 22, 26, 27, 28, 33, and 35, Township 46 South, and Range 28 East.




Figure 1: General Location of and Directions to Pepper Ranch Preserve.

1.4 Regional Significance of the Pepper Ranch Preserve

Ecosystem services such as the protection of water resources, flood control, maintenance of nutrient cycles, preservation of biological diversity, carbon sequestration, and the availability of recreational lands are imperative for the well-being of the citizens of Collier County and may be achieved through the preservation of natural areas. As of February 20173, April 2008 approximately 686% (over 880,980 880,980 acres) of all land in Collier County were protected in conservation areas (Figure 2) and managed by private, local, state and federal agencies (FNAI 2008). Collier County's Conservation Collier Program manages the 2,510.01-acre Pepper Ranch Preserve; it contains upland mixed forest, strand swamp, slough, prairie hammock, pine flatwoods, improved pasture, dry prairie, freshwater marshes, bottomland forest, and wet prairie.

Formatted: Underline

Collier County Preserves, Conservation Easements, **Designated State and Federal Land** Caracara Prairie Preserve - 368 ac Pepper Ranch Preserve - 2,512 ac Red Root Preserve - 9 ac Panther Walk Preserve - 9.5 ac Railhead Scrub Preserve - 130 ac Red Maple Swamp Preserve - 200 ac Winchester Head - 80 ac Lee County Collier County Wet Woods OLWELL RD Preserve - 28 ac MINDKALEE RD Camp Keais Strand Project - 27 ac Alligator Flag Preserve - 20 ac Cocohatchee Rivers Road Preserve - 78 ac Creek Preserve - 4 ac Logan Woods Preserve - 6 ac Freedom Park - 12.5 ac INTERSTATE 75 DAVIS BLV D-Gordon River Greenway Preserve - 41 ac Shell Island Preserve - 80 ac McIlvane Marsh Project - 300 ac Otter Mound Preserve - 3 ac 8 Legend 4 Miles CONSERVATION State and Federal Conservation Land Data Source:Parcels - Collier County Property Apprais Created By: Conservation Collier CS ©\Conservation Collier\maps\Acquired properties\ Acquired_Map_July_2017_2.mxd and .jpg Mitigation / Conservation Easements Collier County

Figure 2: Conserved Lands in Collier County, Florida Including Lands Owned by Conservation Collier.

1.5 Nearby Public Lands and Designated Water Resources

Pepper Ranch Preserve shares its western and a portion of its northern boundary with the Corkscrew Regional Ecosystem Watershed or CREW Trust conservation lands. The closest Conservation Collier Program property to Pepper Ranch Preserve is the Caracara Prairie Preserve, which is approximately 1.23 miles directly to the west. Other preserves, in order of increasing distance, are provided in Table 2. Figure 3 shows the locations of these preserves.

Table 2: Public Lands Located near the Pepper Ranch Preserve				
Name	Distance (miles)	Direction	Туре	
CREW	0.00	W	State	
Caracara Prairie Preserve	1.23	W	Conservation Collier	
Corkscrew Swamp Sanctuary	2.03	SW	National Audubon Society	
Red RootLimpkin Marsh Preserve	5.26	S/SW	Conservation Collier	
Imperial Marsh Preserve	5.63	SE	Lee County Conservation 20/20	




Figure 3: Preserves and Protected Lands in the Vicinity of Pepper Ranch Preserve

1.6 Public Involvement

Community involvement in the review of this management plan and all future updates will be coordinated through the general public via public meetings. The community at large will be contacted through direct mailing notices to residents, other preserve managers and businesses within 1,500 feet of the preserve boundaries. Official public notices will be posted on the County website. Staff will seek to coordinate management actions, such as exotic removal and prescribed fires with managers/owners of adjoining public and private lands.

The off-road cycling group called of the Florida Mudcutters which represent southwest Florida have been active partners since May 2012. They have volunteered hundreds of hours and haveto help developed and maintained biking trails in designated places along the western portion of throughout the preserve.

The Caloosa Saddle Club, a local horseback riding group, has also expressed interest in bringing groups to the Preserve to ride on designated trails. Both groups—have gaveiven County staff input on the conceptual plan for the existing trails.

Several <u>Eagle Scouts and</u> Boy Scouts <u>Groups</u> have volunteered to assist in projects to improve the preserve and trail systems such as building picnic tables, <u>kiosks</u>, <u>hitching post</u>, <u>campground design and development and campground fire rings</u>, marking trails and installing bat boxes.

2.0 Natural Resources

2.1 Physiography

Pepper Ranch Preserve lies within the Floridian section of the Coastal Plain. The Coastal Plain extends from New Jersey to Texas and was formed mainly from sedimentary rocks deposited in marine environments (U.S. Geological Survey, USGS 2004).

2.1.1 Topography and Geomorphology

The site is located in the Southwestern Slope region of the South Florida Water Management District (SFWMD). According to the Florida Geographic Data Layer (FGDL), taken from the USGS Quadrangle Map, the topography of the area is relatively level with an average elevation of twenty feet above sea level and slopes gently southwestward toward the Gulf of Mexico. Surface water percolates directly through the pervious ground or it collects in natural depressions and man made ditches onsite. In natural areas, when the ground is completely saturated the accumulated surface water will drain offsite through sheet flow.

2.1.2 Geology

The geology of northern Collier County, where the Pepper Ranch Preserve is located, is characterized by complex sequences of interbeded sands, clays, and limestone. Closest to the surface is the Holocene aged Pamlico Sand Formation, approximately ten feet thick and composed primarily of unconsolidated quartz sand and some silt. The Pamlico Sand unconformably overlies the Pleistocene aged Fort Thompson and Caloosahatchee Formations, which vary from a few feet to more than twenty feet in thickness and are characterized by shelly and sandy limestone with vugs and solution cavities (Miller 1986).

Below the Fort Thompson and Caloosahatchee Formations are the Ochopee and Buckingham Members of the Pliocene aged Tamiami Formation, which are at least 200 feet thick in the surrounding areas (Oaks & Dunbar 1974). The Ochopee Limestone unconformably overlies the Buckingham Limestone and/or the equivalent Cape Coral Clay. This unconformity marks the bottom of the surficial aquifer separating it from the brackish underlying aquifer below. Then the Hawthorn Formation, rich in phosphate and other heavy minerals (Scott 1988), overlies the Oligocene age Suwannee Limestone and Eocene age Ocala Limestone that form the Floridan Aquifer System in Southwestern Florida. The Pepper Ranch Preserve is located within the Southwestern Slope. Geologically, this is the dominant feature of Collier County (Campbell 1990). Figure 4 provides a current aerial view of the Pepper Ranch Preserve.




Figure 4: Aerial View of the Pepper Ranch Preserve

2.1.3 Soils

Mapped soils on this parcel were identified by the Natural Resource Conservation Services (NRCS) as Riviera Copeland fine sand, Oldsmar fine sand, Riviera fine sand, Ft. Drum and Malabar, Chobee Winder and Gator soils, Boca Riviera and Copeland depressional, Tuscawilla fine sand, Winder Riviera Chobee soils depressional, and Pennsuco silt loam (Figure 5).




Figure 5: Soil Units on the Pepper Ranch Preserve

The following soils descriptions comprise the six hydric or depressional soils at Pepper Ranch Preserve. Winder Riviera Chobee soils underlie 17.6% of the Pepper Ranch Preserve and are very poorly drained soils, or depressional soils; they are typical of marshes. These soils under natural conditions remain ponded, i.e. have standing water, for 6 months or more during most years. Examples of natural vegetation found on these soils include: sawgrass, maidencane, pickerelweed, fireflag, willow, and other wetland plants (Liudahl et al. 1990). Riviera Copeland fine sand, which underlies 13.3% of the Pepper Ranch Preserve, is another hydric or depressional soil found on the preserve. They are a poorly drained soil and are typical of sloughs and cypress swamps. During times of high rainfall, the soils are covered by shallow, slowly moving water for about 7 days. Otherwise during most of the wet season, under natural conditions, the seasonal high water table is within a depth of 12 inches for 3-6 months; and for the remainder of the year the water table is below a depth of 12 inches receding to 40 inches or below during extended dry periods (Liudahl et al. 1990). Chobee Winder and Gator soils comprise 7.4% of Pepper Ranch Preserve. They are very poorly drained soils found in depressions and marshes. Under natural conditions these soils are ponded for 6 months or more of the year, for most years. The water table recedes to within 12 inches the remainder of the year and down to 12-40 inches during extended dry periods. Natural vegetation consists of pickerelweed, maidencane, rushes, fireflag, sawgrass, willow, and a few cypress trees (Liudahl et al. 1990). Pennsuco silt loam is a poorly drained soil

found on low prairies and it comprises 4.1% of the soils found at Pepper Ranch Preserve. Under natural conditions, the seasonal high water table is within a depth of 12 inches for 4-6 months during most years. A few inches of water is above the surface during extremely wet periods. Natural vegetation typically consists of sawgrass, reeds, scattered areas of cypress, maidencane, needlegrass, sedges, wax myrtle, and other wetland plants (Liudahl et al. 1990). The Boca, Riviera and Copeland map unit is a hydric soil that comprises 2.8% of the soils found at Pepper Ranch Preserve. It is found in depressions, cypress swamps, and marshes. Under natural conditions, these soils are ponded for 6 months or more each year. During the remainder of the year the water table is within a depth of 12 inches, and it recedes to a depth of 12-40 inches during extended dry periods. The natural vegetation consists mostly of cypress, pickerelweed, rushes, fireflag, sawgrass, and willow (Liudahl et al. 1990). Only a small percentage of the soils at Pepper Ranch Preserve are Riviera fine sand (0.1%), which is a poorly drained soil found in sloughs and broad, poorly defined drainageways. Under natural conditions, the seasonal high water table is within a depth of 12 inches for 3-6 months during most years. During the other months, the water table is below a depth of 12 inches, and it recedes to a depth of more than 40 inches during extended dry periods. During periods of high rainfall, the soil is covered by shallow, slowly moving water for about 7 days. Natural vegetation found on these soils consists of scattered areas of South Florida slash pine, cypress, cabbage palm, wax myrtle, sand cordgrass, gulf muhly, blue maidencane. South Florida bluestem, and chalky bluestem (Liudahl et al. 1990).

The following soils descriptions comprise the three non-hydric soils found at Pepper Ranch Preserve. Tuscawilla fine sand underlies 30.5% of the Pepper Ranch Preserve and is a non-hydric or non-wetland soil association found in areas typical of flatwoods and hammocks. Under natural conditions, the seasonal high water table is at a depth of 6-18 inches for 1-6 months during most years. The remainder of the year the water table is below 18 inches. During times of drought, the water table can recede to a depth of greater than 40 inches. Natural vegetation consists of oak, cabbage palm, red maple, red bay, South Florida slash pine, wax myrtle, maidencane, and chalky bluestem (Liudahl et al. 1990). Oldsmar fine sand underlies 18.2% of the Pepper Ranch Preserve and is also a non-hydric soil. Oldsmar fine sand is a nearly level and poorly drained soil found in pine flatwoods. During extended dry periods, the water table may recede to a depth of 40+ inches, but under natural conditions, the seasonal high water table is between a depth of 6-18 inches. Flora typically associated with this soil type includes South Florida slash pine, cabbage palm, saw palmetto and wax myrtle (Liudahl et al. 1990). Fort Drum and Malabar fine sands are non-hydric soils typically found on ridges adjacent to sloughs. These soils comprise 4.6% of the soils at Pepper Ranch Preserve. Under natural conditions, the seasonal high water table is at a depth of 6-18 inches for 1-6 months during most years. During the remainder of the year, the water table is below a depth of 18 inches, and it recedes to a depth of more than 40 inches during extended dry periods. Natural vegetation found on these soils is generally South Florida slash pine, saw palmetto, live oak, cabbage palm, wax myrtle, chalky bluestem, creeping bluestem, low panicum and pineland threeawn (Liudahl et al. 1990).

2.1.4 Hydrology/Water Management

Near the surface, the aquifer is highly permeable and the groundwater flows toward the west. However, permeability decreases downward from a porous limestone into poorly indurated sandstone cemented by micrite. The aquifer grades from freshwater downward into brackish water due to the proximity of the Gulf of Mexico to the west and the brackish water in the intermediate aquifer made primarily of Miocene aged sediments. Below that, the Hawthorne formation typically marks the upper boundary of the Floridian aquifer, which is contained within the underlying Oligocene age Suwannee Limestone (Lodge 2005).

There are numerous ditches and berms at Pepper Ranch Preserve that are associated with the former agricultural activities. There are also elevated roads with associated ditches that run through the center of the preserve, initially in an east-west direction, then running north-south all the way to the southwestern portion of the preserve (see Figure 6 for land use and cover map). These ditches and berms are likely affecting the hydrology of the wetlands in which they connect.

The Surficial Aquifer is an aquifer close to the surface and unconfined, typically associated with the groundwater table. This aquifer is generally limited to smaller uses such as household or small agricultural uses. The Lower Tamiami aquifer is below this aquifer and is recognized as being useful for long-term water needs. According to the SFWMD's technical publication 95-02 (Fairbank & Hohner 1995), the Surficial Aquifer recharge capacity on the Pepper Ranch Preserve is moderate at 43 to 56 inches annually, with parts of the eastern portion of the preserve exhibiting a recharge capacity of less than 43 inches annually. The Lower Tamiami Aquifer recharge capacity on the preserve is relatively low ranging from less than 7 inches and up to 14 inches annually.

2.2 Climate

The Pepper Ranch Preserve is located in an area of Florida where humid subtropical and tropical savanna climatic patterns overlap, with temperatures moderated by winds from the Gulf of Mexico and the Atlantic Ocean. Sharply delineated wet and dry seasons and average monthly temperatures greater than 64° Fahrenheit characterize a tropical savanna climate. Monthly rainfalls may exceed ten inches during the wet season. On the other hand, humid subtropical climates typically show less extreme rainfall fluctuations between wet and dry seasons and average monthly temperatures are less than 64° Fahrenheit in some months.

The average annual temperature for the coastal portion of Collier County is approximately 75° Fahrenheit. The warmest months are usually July and August. The humidity is high during these months but frequent afternoon thunderstorms prevent excessively high temperatures. Two-thirds of the annual rainfall occurs in the wet season from May to October. Thunderstorms are frequent during the wet season, occurring every two out of three days between June and September. Rainfall records for the area indicate that there is not significant variation in the annual rainfall throughout much of the county; however, large variations often occur during a single year. The Atlantic hurricane season extends from June through November with peak activity

occurring in September and October when ocean temperatures are highest.

2.3 Natural Plant Communities

A plant community refers to the suite of floristic species that form the natural vegetation of any place. In addition to anthropogenic influences, the combination of factors such as geology, topography, hydrology, underlying soils and climate determine the types of plants found in an area. These plants, in turn determine the animal species that may be found there. The description or classifications of these floral communities differ by agency and are based on an agency's goals and objectives for identifying plant communities. As some categorizations are broad (e.g., forest) while others are specific (e.g., mesic pine flatwoods), determining how each organization classifies a community may be difficult. The plant communities observed on the Pepper Ranch Preserve are presented using the Florida Land Use, Cover and Forms Classification System (FLUCFCS) created by the Florida Department of Transportation (1999). This system classifies all land uses including plant communities. These classifications were then translated to the Florida Natural Areas Inventory (FNAI) classifications. The Guide to the Natural Communities of Florida (1990) was utilized to convert from FLUCFCS to FNAI designations.

In the fall of 2009 Johnson Engineering ecologists mapped the vegetation communities and other land uses found on the Pepper Ranch Preserve using the FLUCFCS designations. There are 25 distinct plant communities/land uses on the preserve. Some of these land uses are further described as disturbed and/or by the level of invasive exotic plants they exhibit. The number 9 qualifier in the FLUCFCS code represents a disturbance in the plant community, generally due to a hydrologic impairment and in some cases the disturbance is from a cleared understory. The letter E qualifier represents the level of invasive exotic vegetation present by percent cover. Table 3 summarizes the plant communities mapped for the Pepper Ranch Preserve in 2009. The table also provides a brief description of each FLUCFCS code. Figure 6 visually depicts these land cover designations from 2009. Due to the size of Pepper Ranch Preserve, a 2'x3' map of the land cover designations for 2009 is provided in Appendix 2.

Table 3: Extent of Florida Land Use, Cover and Forms Classification System (FLUCFCS) Designations from 2009 on the Pepper Ranch Preserve			
FLUCFCS Code	Description	Wetland Status	Acreage
1641	Oil field	N	5.63
211	Improved pasture	N	619.26
2111	Cattle pen	N	0.38
3109	Upland prairie, disturbed	N	3.73
310E1	Upland prairie, disturbed, exotics 5-24%	N	37.35
310E2	Upland prairie, disturbed, exotics 25-49%	N	4.01

Table 3: Extent of Florida Land Use, Cover and Forms Classification System (FLUCFCS)
Designations from 2009 on the Pepper Ranch Preserve (continued)

FLUCFCS Code	Description	Wetland Status	Acreage
3209	Upland shrub, disturbed	N	3.53
411	Pine flatwoods	N	149.95
4119	Pine flatwoods, disturbed	N	40.80
4119E1	Pine flatwoods, disturbed, exotics 5-24%	N	1.44
4119E2	Pine flatwoods, disturbed, exotics 25-49%	N	27.45
4119E4	Pine flatwoods, exotics 75-100%	N	3.69
422	Brazilian pepper, non-hydric	N	1.15
427/428	Oak/Cabbage palm	N	11.22
427E1	Oak, exotics 5-24%	N	1.57
428E1	Cabbage palm, exotics 5-24%	N	3.82
428E3	Cabbage palm, exotics 50-74%	N	4.31
434	Oak, slash pine, cabbage palm	N	270.92
4349	Oak, slash pine, cabbage palm, disturbed	N	1.99
4349E1	Oak, slash pine, cabbage palm, disturbed, exotics 5-24%	N	149.74
4349E2	Oak, slash pine, cabbage palm, disturbed, exotics 25-49%	N	8.86
434B	Oak, slash pine, cabbage palm, burned	N	34.65
437	Australian pine	N	1.15
743	Spoil	N	1.24
743E4	Spoil, exotics 75-100%	N	9.49
8145	Shell road, graded and drained	N	20.63
8146	Primitive trail	N	4.90
512	Ditches	OSW	34.49
512E4	Ditches, exotics 75-100%	OSW	0.66
742	Borrow pond	OSW	1.43
211H	Improved pasture, hydric	W	47.70

Table 3: Extent of Florida Land Use, Cover and Forms Classification System (FLUCFCS)

Designations from 2009 on the Pepper Ranch Preserve (continued)

FLUCFCS Code	FLUCFCS Code	FLUCFCS Code	FLUCFCS Code
6151	Red maple swamp	W	76.56
6152	Pop ash swamp	W	15.05
6162	Pond apple depression	W	0.71
6169E1	Pond apple, laurel oak, cabbage palm, disturbed, exotics 5-24%	W	0.83
6189	Willow/shrub wetland, disturbed	W	5.44
6192	Brazilian pepper, hydric	W	4.26

621	Cypress	W	82.41
624	Cypress, pine, cabbage palm	W	2.51
624E1	Cypress, pine, cabbage palm, exotics 5-24%	W	41.15
630	Wetland forested mix	W	145.46
630E1	Wetland forested mix, exotics 5-24%	W	3.85
631	Shrub wetland	W	243.38
6319E1	Shrub wetland, disturbed, exotics 5-24%	W	12.26
6319E2	Shrub wetland, disturbed, exotics 25-49%	W	0.56
631E1	Shrub wetland, exotics 5-24%	W	1.35
641	Freshwater marsh	W	42.60
6419	Freshwater marsh, disturbed	W	44.83
6419E1	Freshwater marsh, disturbed, exotics 5-24%	W	70.99
6419E2	Freshwater marsh, disturbed, exotics 25-49%	W	26.19
6419E3	Freshwater marsh, disturbed, exotics 50-74%	W	5.92
641E1	Freshwater marsh, exotics 5-24%	W	135.72
641E2	Freshwater marsh, exotics 25-49%	W	26.37
641E3	Freshwater marsh, exotics 50-74%	W	1.27
643	Wet prairie	W	5.21
6439E2	Wet prairie, disturbed, exotics 25-49%	W	7.50
643E1	Wet prairie, disturbed, exotics 5-24%	W	0.49
		Total	2510.01

 $\begin{aligned} W - wetland \\ N - non-wetland \end{aligned}$

OSW – other surface water




Figure 6: Distribution of Natural Communities and other Land Uses on the Pepper Ranch Preserve; 2009 FLUCFCS Layer

The vegetation classification scheme of the FNAI and the Florida Department of Natural Resources (FDNR) (1990) are presented in Table 4. This table is based on the natural plant communities observed on the Pepper Ranch Preserve. The following subsections (2.3.1 - 2.3.6) provide information about the natural plant communities observed on the preserve according to their FNAI designations. Subsection 2.3.7 describes the altered communities found at Pepper Ranch Preserve. Figure 7 visually depicts the FNAI designations for the preserve based on the 2009 field verifications. Due to the size of Pepper Ranch Preserve, a 2'x3' map of the FNAI designations is provided in Appendix 3.

Table 4: Summary of Natural Communities on the Pepper Ranch Preserve						
FNAI Natural Community Type	Global Rank	State Rank	Percent Cover ¹	Comments		
Bottomland forest	G4	S3	9.87%	Also called bottomland hardwoods and mesic hammock		
Depression marsh	G4	S4	14.09%	Also called isolated wetland and ephemeral pond		
Dry prairie	G2	S2	1.94%	Also called palmetto prairie		
Mesic flatwoods	G4	S4	8.97%	Also called pine flatwoods		
Prairie Hammock	G3	S3	0.83%	Also called palm/oak hammock and hydric hammock		
Slough	G3	S3	10.25%			
Strand swamp	G4	S4	3.28%	Also called cypress strand		
Upland mixed forest	G4	S4	18.56%	Also called upland hardwood and mesic hammock		
Wet flatwoods	G4	S4	1.74%	Also called hydric flatwoods		
Wet prairie	G3	S2	0.53%	Also called savannah and coastal prairie		

^{70.06%} of Pepper Ranch Preserve is comprised of natural communities. The remaining 29.94% is comprised of altered communities as described in subsection 2.3.7.

Definition of Global (G) element ranks:

- G2 = Imperiled globally because of rarity (6-20 occurrences or very little remaining area, e.g., <10,000 acres) or because of some factor(s) making it very vulnerable to extinction throughout its range:
- G3 = Either very rare and local throughout its range or found locally (even abundantly at some of its locations) in a restricted range or because of other factors making it vulnerable to extinction throughout its range, 21 to 100 occurrences;
- G4 = Apparently secure globally, though it may be quite rare in parts of its range, especially at the periphery.

Definition of State (S) element ranks:

- S2 = Imperiled in state because of rarity (6-20 occurrences or little remaining area) or because of some factor(s) making it very vulnerable to extinction throughout it range:
- S3 = Rare or uncommon in state (on the order of 21 to 100 occurrences);
- S4 = Apparently secure in state, although it may be rare in some parts of its state range.




Figure 7: Extent of Natural Plant Communities Currently Found on the Pepper Ranch Preserve.

2.3.1 Uplands: Upland Mixed Forest

The upland mixed forest community comprises approximately 18.56% of the Pepper Ranch Preserve, the largest acreage of which is located primarily in the southwestern portion of the preserve but also occurs throughout the preserve in smaller acreages (Figure 7).

Upland mixed forests in south Florida are also known as upland hardwoods, mesic hammocks, prairie hammocks, xeric hammocks, hydric hammocks (FNAI 1990) and mesic temperate hammocks (USFWS 1999). This plant community at Pepper Ranch Preserve is characterized by live oaks (Quercus virginiana) and laurel oaks (Quercus laurifolia) (both hardwood species), south Florida slash pine (Pinus elliottii var. densa), cabbage palms (Sabal palmetto), and some cypress (Taxodium spp.) and strangler figs (Ficus aurea), that together



Upland Mixed Forest Community within Pepper Ranch Preserve Photo by Johnson Engineering, Inc.

generally form a closed canopy. The midcanopy includes myrsine (Myrsine guainensis), dahoon holly (Ilex cassine) and wax myrtle (Myrica cerifera). The plant species found in the understory are mainly native species including blue maindencane (Amphicarpum muhlenbergianum), foxtail grass (Setaria sp.), carpet grasses (Axonopus spp.), slender goldenrod (Euthamia caroliniana), musky mint (Hyptis alata), chocolate weed (Melochia corchorifolia), swamp fern (Blechnum serrulatum), capeweed (Phyla nodiflora), wild coffee (Psychotria nervosa) and varying densities of the invasive exotic torpedo grass (Panicum repens). Originally in theIn—the disturbed portions of this community. Brazilian pepper (Schinus terebinthifolius) compriseds up to 24% of the midcanopy otherwise it is present at less than 5% coverage. In the herbaceous layer of the disturbed areas of this community there wereare significant levels (26-50% coverage) of caesarweed (Urena lobata) and some (1-5% coverage) dogfennel (Eupatorium capillifolium). These areas have been treated several times since acquisition, however a significant seed source exists. Ongoing maintenance is being conducted on an annual or bi-annual basis.

The closed canopy and abundant hardwood mast provided by this plant community attracts a number of wildlife species seeking food, cover, roosting, and nesting sites. Additionally, these areas are ideal stopover areas for migratory passerines. Since these communities occur on relatively well-drained sites, they are also attractive for human habitation and recreational uses. These anthropogenic uses have increased the number of

invasive plant and animal species in these areas and have resulted in degraded hardwood forests throughout the state (USFWS 1999).

2.3.2 Wetlands: Depression Marsh

Depression marsh, also known as freshwater marsh, isolated wetland, or ephemeral pond, comprises approximately 14.09% of the Pepper Ranch. Even though these wetlands are present throughout the preserve, they are more prevalent in the central portion of the preserve and eastward to the property boundary.

Freshwater marshes are often scattered among upland, pine flatwoods communities as is the case at Pepper Ranch Preserve. Only 12% of this plant community at Pepper Ranch

Preserve appears to be hydrologically undisturbed. This determination was made solely by observing the plant composition in the disturbed marshes, which consisted of varying levels of upland and transitional species, nuisance and invasive exotic species, as well as by noting the relatively low water levels or lack there of in comparison to the seemingly undisturbed marshes. Further hydrologic investigations would be necessary to determine if hydrology was the actual cause of the noted disturbance.



Depression Marsh Community within Pepper Ranch Preserve Photo by Johnson Engineering, Inc.

Native plant species found within this plant community included maidencane (Panicum hemitomon), Southeastern sunflower (Helianthus agrestis), blue maidencane, American cupscale (Sacciolepis striata), pickerelweed (Pontederia cordata), alligator flag (Thalia geniculata), swamp fern, sawgrass (Cladium jamaicense), narrowfruit horned beaksedge (Rhyncospera inundata), southern beaksedge (Rhyncospora microcarpa), pale meadowbeauty (Rhexia mariana), bulltongue arrowhead (Sagittaria lancifolia), musky mint, Virginia buttonweed (Diodea virginiana), common buttonbush (Cephalanthus occidentalis), lemon bacopa (Bacopa caroliniana), spikerushes (Eleocharis spp.), American white waterlily (Nymphaea odorata), broomsedge bluestem (Andropogon virginicus), sand cordgrass (Spartina bakeri), redtop panicum (Panicum rigidulum), and corkwood (Stillingia aquatica). The nuisance and invasive exotic plant species observed in this community include torpedo grass, dog fennel, caesarweed, Southern crabgrass (Digitaria ciliaris), tropical soda apple (Solanum viarum), alligator weed (Alternanthera philoxeroides), Brazilian pepper, melaleuca (Melaleuca quinquenervia), Peruvian

primrosewillow (Ludwigia peruviana), and valamuerto (Senna pendula var. glabrata).

In Florida, these marshes are influenced by their subtropical location, fluctuating water levels, frequency and intensity of fire, organic matter accumulation and hard water (Kushlan 1990). These factors, combined with the dominant species found within a marsh, dictate the category within which the marsh is placed. Six major categories of freshwater marshes are recognized in Florida. The marshes in the Pepper Ranch Preserve are generally within the "flag marsh" category. These marshes usually have a moderate (flooded 6 to 9 months) hydroperiod, a moderate (about once every ten years) fire frequency and moderate to high (usually less than one meter deep to over a meter deep) accumulation of organic material (Kushlan 1990).

2.3.3 Wetlands: Slough

Sloughs are generally abundant throughout Florida and at Pepper Ranch Preserve this community comprises 10.25% of the preserve. It is located almost entirely in the western portion of the preserve, and most of this community extends onto the adjacent CREW land.

According to the FNAI Guide to the Natural Communities of Florida (1990), sloughs are characterized as broad shallow channels, inundated with flowing water except during extreme droughts. They are the deepest drainageways within strand swamps and swale systems. The vegetation structure of sloughs is variable but at Pepper Ranch Preserve it is characterized, in general, by Carolina willow (Salix caroliniana), which is dominate, Carolina (pop) ash (Fraxinus caroliniana). Sawgrass, bog hemp (Boehmeria cylindrica) and climbing hempweed (Mikania scandens) were some of the herbaceous species observed in the understory.

The canopies formed in these sloughs, especially in south Florida, are ideal moist, warm habitats for rare and endangered tropical epiphytes. Many Caribbean species that occur in this community are virtually never



Slough Community within Pepper Ranch Preserve Photo by Johnson Engineering, Inc.

encountered in other Florida habitats. Pond apple branches are often heavily loaded with epiphytic orchids, bromeliads, and ferns. Typical animals include ribbon snake,

cottonmouth, opossum, gray squirrel, black bear, raccoon, mink, otter, Florida panther, and white-tailed deer (FNAI 1990).

Sloughs often occur over the lowest part of linear depressions in the underlying limestone bedrock. The peat soils found in sloughs can be destroyed by catastrophic fires that often occur during droughts. The typical hydroperiod in this community is at least 250 days per year. Sloughs are often found in association with cypress swamps and may also occur in floodplain swamps and basin swamps (FNAI 1990).

Sloughs are extremely vulnerable to hydrologic disturbance and must have a reliable, quality water source to persist. The lack of invasive plant species observed in this community at Pepper Ranch Preserve is indicative of a high quality system.

2.3.4 Wetlands: Bottomland Forest

This community at the Pepper Ranch Preserve appears in association with Lake Trafford along the southern perimeter of the preserve, and with the large slough occurring on the western portion of the preserve. This natural community covers 9.87% of the preserve.

Bottomland forest is characterized as a low-lying, closed-canopy forest of tall, straight trees with either a dense shrubby understory and little ground cover, or an open understory and ground cover of ferns, herbs, and grasses (FNAI 1990). At Pepper Ranch Preserve the latter is most often observed, with red maple (Acer rubrum) as the dominate canopy tree, some buttonbush in the midcanopy and sawgrass, alligator flag, swamp fern, and cinnamon fern (Osmunda cinnamomea) the understory.

The canopy of these forests is dense and closed, except during winter in areas where deciduous trees predominate, as in Pepper Ranch Preserve. The air movement and light penetration are thus generally low, making the humidity high and relatively constant. Because of these characteristics, bottomland forests rarely burn. This is also a very stable community that requires a hundred years or more to mature. Nearly all



Bottomland Forest Community within Pepper Ranch Preserve Photo by Johnson Engineering, Inc.

bottomland forests in Florida have been logged, which often leaves long-lasting scars from soil disturbance (FNAI 1990).

2.3.5 Uplands: Mesic Flatwoods

The pine flatwoods community comprises approximately 8.97% of the Pepper Ranch Preserve. This plant community is located predominately in the central and eastern portions of the preserve. Pine flatwoods are one of the most wide-ranging terrestrial plant communities in Florida and consequently one of the most influenced by anthropogenic activities (Abrahamson & Hartnett 1990). Fire strongly influences the community structure and composition of this community. The term pine flatwoods is a general categorization of areas that are dominated by various species of pine (*Pinus* spp.) trees. Pine flatwoods may be found in mesic flatlands where the landscape is made up of flat, moderately well drained sandy substrates with a mixture of organic material, often with an underlying hard pan layer. An open canopy forest of widely spaced pine trees with little or no understory but a dense ground cover of herbs and shrubs characterize natural, mesic flatwoods that have been burned regularly (FNAI 1990).



Mesic Flatwoods Community within Pepper Ranch Preserve Photo by Steven W. Woodmansee

Department The U.S. Agriculture (USDA) NRCS classification system refers to these areas as South Florida South Florida flatwoods. flatwoods are typically savannas, a type of plant community intermediate between forest and grassland. Mesic pine flatwoods are also called mesic flatwoods, pine savanna, cabbage palm savanna, and pine barrens. The flatwoods at Pepper Ranch are characterized by a south Florida slash pine dominate canopy with some live oaks, wax myrtle and saw palmetto (Serenoa repens) in the

subcanopy, and a myriad of herbs and forbs forming the ground cover, such as: swamp fern, grape vine (*Vitis rotundifolia*), American beautyberry (*Callicarpa americana*), tall elephant's foot (*Elephantopus elatus*), greenbrier (*Smilax* sp.), caesarweed, and ticktrefoil (*Desmodium* sp.).

Mesic flatwoods provide essential forested habitat for a variety of wildlife species including Neotropical migratory birds, wide-ranging large carnivores, mid-sized carnivores, ground-nesting vertebrates, tree-cavity dependent species, tree-nesting species and non-aquatic plant life. "At the current rate of habitat conversion, the mesic pine flatwoods, once the most abundant upland habitat in South Florida, is in danger of becoming one of the rarest habitats in South Florida" (USFWS 1999c).

2.3.6 Other Natural Communities

All other natural communities (strand swamp, dry prairie, wet flatwoods, prairie hammock, and wet prairie) at the Pepper Ranch Preserve collectively cover less than 9% of the preserve.

Strand swamp is most commonly known as cypress swamp and it is strictly found in the western portion of Pepper Ranch Preserve in association with the slough natural community. The typical vegetation found in this community at Pepper Ranch Preserve includes: cypress (*Taxodium ascendens*) in the canopy, Carolina (pop) ash in the midcanopy, and the understory is mainly open water with some alligator flag, pickerelweed, and sawgrass.

The dry prairie community at Pepper Ranch Preserve is located in the extreme southwest and eastern portions of the preserve. Portions of this community exhibit no canopy, but where a canopy is present it is at less than 10% coverage and is made up of south Florida slash pine and cabbage palms. The midcanopy appears to have



Strand Swamp Community within Pepper Ranch Preserve Photo by Johnson Engineering, Inc.



Dry Prairie Community within Pepper Ranch Preserve Photo by Johnson Engineering, Inc.

been cleared at one time and now consists mainly of saw palmetto and wax myrtle. The understory is dominated by bahiagrass (Paspalum notatum), and other grasses and herbs present including: blue maidencane, torpedo grass, knotroot foxtail (Setaria parviflora), carpetgrass (Axonopus furcatus), chocolate weed, tall elephant's foot, ticktrefoil, musky mint, netted pawpaw (Asimina reticulata), slender goldenrod, wire grass (Aristida stricta), and a variety of sedges. The disturbed portions of this community located in the eastern part of the preserve exhibit less native plant diversity, which has been replaced by

cogongrass (*Imperata cylindrica*), smutgrass (*Sporobolus indicus*), Peruvian primrose willow, caesarweed, and Brazilian pepper. The disturbance to the portions of this community located in the southwestern part of the preserve appears to be due to a cleared canopy. The invasive plants in these areas have been treated several times since acquisition, however a significant seed source exists. Ongoing maintenance is being conducted on an annual or bi-annual basis.

The wet flatwoods of Pepper Ranch Preserve are exclusively found in the western portion of the preserve and they exhibit cypress, south Florida slash pine and cabbage palms in the canopy, little to no midcanopy and an understory similar to the adjacent strand swamp community.

The prairie hammocks at Pepper Ranch Preserve are an upland community comprised of predominately live oaks and cabbage palms in the canopy and a midcanopy and understory similar to the upland mixed forest as described above.



Prairie Hammock Community within Pepper Ranch Preserve Photo by Johnson Engineering, Inc.

following plant species: blue maidencane, sand cordgrass, corkwood, broomsedge bluestem, haspan flatsedge (Cyperus haspan), spadeleaf (Centella asiatica), cypress witchgrass (Dichanthelium unciphyllum), ensifolium var. southern umbrellasedge (Fuirena scirpoidea), maidencane, narrowfruit horned beaksedge, narrowleaf blueeyed grass (Sisyrinchium angustifolium). Portions of this community also have the invasive exotic torpedo grass, up to 50% coverage, and the remaining portions are free of invasive exotic vegetation. These invasive plant areas have been treated several times since acquisition, however a significant seed source exists. maintenance is being conducted on an annual or biannual basis.



Wet Flatwoods Community within Pepper Ranch Preserve Photo by Steven W. Woodmansee

There are only four small areas of wet prairie at Pepper Ranch Preserve located in the southwestern portion of the preserve and they exhibit the



Wet Prairie Community within Pepper Ranch Preserve Photo by Johnson Engineering, Inc.

2.3.7 Altered Communities

The most common community at Pepper Ranch Preserve is the improved pastures, which is an altered community; they comprise 26.57% of the preserve. The majority of the pastures exhibit upland grass and forbs species dominated by bahiagrass with a mixture of the following species: limpograss (*Hemarthria altissima*), ragweed (*Ambrosia*



Improved Pastures within Pepper Ranch Preserve Photo taken by Johnson Engineering, Inc.

artemisiifolia), dogfennel, smutgrass, bushy bluestem (Andropogon glomeratus var. hirsutior), tick-trefoil, capeweed, creeping woodsorrel (Oxalis corniculata), rabbitbells (Crotalaria rotundifolia), flatsedges, torpedo grass, purple thistle (Cirsium horridulum), knotroot foxtail, big carpetgrass, buttonweed woodland false (Spermacoce assurgens), and crabgrass. Most of the pastures at Pepper Ranch Preserve are rimmed with large Brazilian pepper trees associated with ditch/berm and fence lines. In the far western portion of the preserve there is a hydric pasture that exhibits more

wetland species than the other pastures and the underlying soils are hydric soils. There is a midcanopy in the hydric pasture of pop ash, Brazilian pepper, and Carolina willow. The herbaceous layer consists of southeastern sunflower, torpedo grass, bushy bluestem, spadeleaf, Virginia buttonweed, blue mistflower (*Conoclinium coelestinum*), musky mint, and southern beaksedge.

Three oil fields are located adjacent to the hydric pasture at Pepper Ranch Preserve, two of which are currently in operation. An elevated, graded shell road traverses the preserve and provides access to the oil fields in the western portion of the preserve. There are also numerous primitive roads and trails that provide access to the pastures and to the lodge; the latter is located in the southeastern portion of the preserve. Ditches are typically associated with the roads, trails, oil fields and pastures of the preserve. A few borrow ponds are scattered throughout the preserve, usually located within pastures to provide water for the cattle. These borrow ponds typically have spoil piles, some of which are covered with Brazilian pepper.

There are only a few areas where exotics have formed a monoculture. The largest of these areas is located at the southern end of a large depression marsh in the north portion of the preserve where there is a hydric Brazilian pepper monoculture (4.26 acres). In the southeastern portion of the preserve there wereare two small pockets of mature suckering Australian pines (*Casuarina glauca*) totaling 1.15 acres. These areas have been treated several times since acquisition, however a significant seed source exists. Ongoing maintenance is being conducted on an annual or bi-annual basis. This area has been treated and ongoing maintenance is being conducted on an annual or bi-annual basis.

2.4 Native Plant and Animal Species

The Pepper Ranch Preserve is composed of several upland and wetland natural communities as well as altered communities such as the dominant feature, the improved pastures. This section discusses the flora and fauna observed within these communities and the next section (2.5) discusses all listed species in greater detail.

2.4.1 Plant Species

To date, 416 plant species have been recorded at the preserve (Appendix 4). A comprehensive plant survey was conducted in September 2009 by botanist Steven W. Woodmansee of Pro Native Consulting. An additional survey of Pepper Ranch Preserve was conducted in May of 2010 to capture species in bloom that might have been missed during the fall survey. Of these 416 species, 334 (80%) are native to Florida and 82 are non-native (20%). Of the 82 non-native species, 32 are listed on Florida Exotic Pest Plant Council's (FLEPPC) 2017.09 List of Invasive Plant Species (243 Category I and 8 nine Category II).

2.4.2 Animal Species

When this original plan was written, Due to the there were limited surveys conducted specifically for the occurrence of animal species (in contrast to plants) and the lack of onsite staffing, little wasis recorded for actual occurrences of animals at the Pepper Ranch Preserve. Occurrences of fauna at the preserve wereare based on direct visual and aural observations made by staff, Johnson Engineering ecologists, and Pro Native Consulting biologist during site visits or evidence of activity such as spoor, scat, or burrows, and from the site information available in documents such as the site's initial criteria screening report, the property's interim management plan and anecdotal information from persons with knowledge of the site. Since acquisition, numerous wildlife cameras have been deployed throughout the preserve to give us a more complete list. Table 5 provides a comprehensive list of animals, both native and non-native, recorded on the Pepper Ranch Preserve thus far.

Table 5: Faunal Species Observed at Pepper Ranch Preserve					
Common Name	Scientific Name	Protection Status			
American Bittern	Botaurus lentiginosus				
American Kestrel	Falco sparverius				
American Redstart	Setophaga ruticilla	·			
American Robin	Turdus migratorius				
Anhinga	Anhinga anhinga				
Audubon's Crested Caracara	Polyborus plancus audubonii	T (FWC, USFWS)			
Bald Eagle	Haliaeetus leucocephalus	T (FWC) Delisted (USFWS & FWC)			
Barred Owl	Strix varia				
Belted Kingfisher	Megaceryle alcyon				
Black Vulture	Coragyps atratus				

Formatted: Highlight
Formatted: Highlight
Formatted: Highlight
Formatted: Highlight

Blue-gray Gnatcatcher	Polioptula caerulea				
Table 5: Faunal Species Observed at Pepper Ranch Preserve (continued)					
Common Name	Scientific Name	Protection Status			
Blue Jay	Cyanocitta cristata				
Brown Thrasher	Toxostoma rufum				
Carolina Wren	Thryothorus ludovicianus				
Cattle Egret	Bubulcus ibis				
Common Bobwhite	Colinus virginianus				
Common Grackle	Quiscalus quiscula				
Common Moorhen	Gallinula chloropus				
Downy Woodpecker	Picoides pubescens				
Eastern Meadowlark	Sturnella magna				
Florida Sandhill Crane	Grus canadensis pratensis	T (FWC)			
Black-Bellied Whistling Duck	Dendrocygna autumnalis				
Gray Catbird	Dumetella carolinensis				
Great Blue Heron	Ardea herodias				
Great Crested Flycatcher	Myiarchus crinitus				
Great Egret	Ardea alba				
Green Heron	Butorides virescens				
Limpkin	Aramus guarauna	SSC(1)(FWC)			
Little Blue Heron	Egretta caerulea	<u>TSSC (1,4)</u> (FWC)			
Loggerhead Shrike	Lanius ludovicianus				
Mourning Dove	Zenaida macroura				
Northern Cardinal	Cardinalis cardinalis				
Northern Mockingbird	Mimus polyglottos				
Osprey	Pandion haliaetus				
Pileated Woodpecker	Dryocopus pileatus				
Pine Warbler	Dendroica pinus				
Red-bellied Woodpecker	Melanerpes carolinus				
Red-shouldered Hawk	Buteo lineatus				
Red-winged Blackbird	Agelaius phoeniceus				
Roseate Spoonbill	Ajaia ajaja	<u>TSSC (1,4)</u> (FWC)			
Snowy Egret	Egretta thula	SSC (1)(FWC)			
Swallow-tailed Kite	Elanoides forficatus				
Tree Swallow	Tachycineta bicolor				
Tricolored Heron	Egretta tricolor	<u>TSSC (1,4)</u> (FWC)			
Turkey Vulture	Cathartes aura				
White-eyed Vireo	Vireo griseus				
White Ibis	Eudocimus albus	SSC(2)(FWC)			
Wild Turkey	Meleagris gallopavo				
Wood Stork	Mycteria americana	TE-(FWC), TE (USFWS)			

Yellow-crowned Night-Heron	Nyctanassa violacea					
Yellow-rumped Warbler	Dendroica coronata					
Table 5: Faunal Species Observed at Pepper Ranch Preserve (continued)						
Common Name	Scientific Name	Protection Status				
Armadillo	Dasypus novemcinctus					
Big Cypress Fox Squirrel	Sciurus niger avicennia	T (FWC) _not observed by staff				
Bobcat	Lynx rufus					
Coyote	Canis latrans					
Eastern Cottontail Rabbit	Sylvilagus floridanus					
Feral Hog	Sus scrofa					
Florida Black Bear	Ursus americanus floridanus	T (FWC)				
Florida Panther	Puma concolor coryi	E (FWC); E (USFWS)				
Grey fox	<u>Urocyon cinereoargenteus</u>					
Grev Squirrel	Sciurus carolinensis					
Opossum	Didelphis virginiana					
Raccoon	Procyon lotor					
River otter	Lontra canadensis					
Round-tailed Muskrat	Neofiber alleni					
White-tailed Deer	Odocoileus virginianus					
American Alligator	Alligator mississippiensis	TSSC (FWC); T (USFWS) ¹				
Black Racer	Coluber constrictor priapus					
Brown Anole	Anolis sagrei					
Brown w₩atersnake	<u>Nerodia taxispilota</u>					
Coral Snake	Micrurus fulvius					
Cottonmouth Snake	Agkistrodon piscivorus					
Diamondback Rattlesnake	Crotalus adamanteus					
Florida Box Turtle	Terrapene carolina bauri					
Florida Softshell	Apalone ferox					
Gopher Tortoise	Gopherus polyphemus	T (FWC)				
Green Anole	Anolis carolinensis					
Pigmy Rattlesnake	Sistrurus miliarius					
Florida Softshell Snapping Turtle	Apalone feroxChelydra serpentina					
Three-striped Mud Turtle	Kinosternon bauri					
1	I	1				

Eastern Narrowmouth Toad
List of Abbreviations:

Florida Cricket Frog

Yellow Rat Snake

Pig Frog

FWC = Florida Fish and Wildlife Conservation Commission

Rana grylio

Formatted: Left

Formatted: Font: (Default) Times New Roman, 9 pt

Formatted: Font: (Default) Times New Roman, 9 pt

Elaphe obsoleta quadrivittata

Acris gryllus dorsalis

Gastrophryne carolinensis

USFWS = United States Fish and Wildlife Service

E = Endangered

T = Threatened

SSC = Species of Special Concern

Reasons for SSC listings prior to January 1, 2001 as indicated by number in parenthesis (1)—Has a significant vulnerability to habitat modification, environmental alteration, human disturbance, or human exploitation which, in the foreseeable future, may result in its becoming a threatened species unless appropriate protective or management techniques are initiated or maintained

- (2) May already meet certain criteria for designation as a threatened species but for which conclusive data are limited or lacking
- (1) (4) Has not significantly recovered from past population depletion

The Florida Breeding Bird Atlas (FWC 2003) lists 49 avian species that have been recorded as confirmed, probable, or possible breeding in the vicinity of the site (Table 6). The Breeding Bird Atlas documents breeding distributions of all bird species in Florida between 1986 and 1991. Due to the size and diversity of natural communities found at Pepper Ranch Preserve, it is likely several of these species may breed at the preserve.

Other wildlife species that have not yet been recorded undoubtedly occur at the Pepper Ranch Preserve. During migration periods, transient bird species would be expected to utilize this area for short periods of time. The developed character of the adjacent areas may inhibit transient use by many mammal, reptile, and amphibian species, thus limiting the utilization of the preserve to resident individuals or inhibiting the dispersal of many species to and from the preserve.

Table 6: Breeding Bird Species Recorded in the Corkscrew and Immokalee

Quadrangles Encompassing the Pepper Ranch Preserve Common Name Scientific Name **Common Name** Scientific Name Green Heron Red-cockaded Woodpecker Butorides virescens Picoides borealis Yellow-crowned Nyctanassa violacea Northern Flicker Colaptes auratus Night-Heron Wood Duck Aix sponsa Pileated Woodpecker Dryocopus pileatus Mottled Duck Anas fulvigula Great Crested Flycatcher Myiarchus crinitus Swallow-tailed Kite Elanoides forficatus Loggerhead Shrike Lanius ludovicianus Red-shouldered Hawk White-eyed Vireo Buteo lineatus Vireo griseus Northern Bobwhite Colinus virginianus Blue Jav Cyanocitta cristata King Rail Fish Crow Rallus elegans Corvus ossifragus Common Moorhen Gallinula chloropus Purple Martin Progne subis Limpkin Northern Rough-winged Stelgidopteryx serripennis Aramus guarauna Swallow Killdeer Charadrius vociferus Tufted Titmouse Baeolophis bicolor Mourning Dove Zenaida macroura Carolina Wren Thryothorus ludovicianus Common Ground-Dove Columbina passerina Blue-gray Gnatcatcher Polioptilia caerulea *Rose-ringed Parakeet Psittacula krameri Northern Mockingbird Mimus polyglottos

Brown Thrasher

Northern Parula

Prairie Warbler

Pine Warbler

Toxostoma rufum

Dendroica pinus

Parula americana

Dendroica discolor

Formatted: No bullets or numbering

Formatted: Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Tab after: 0.54" + Indent at: 0.54"

Coccyzus americanus

Megascops asio

Bubo virginianus

Tyto alba

Yellow-billed Cuckoo

Eastern Screech-Owl

Great Horned Owl

Barn Owl

Barred Owl	Strix varia	Common Yellowthroat	Geothlypis trichas
Common Nighthawk	Chordeiles minor	Eastern Towhee	Pipilo erythrophthalmus
Chuck-will's-widow	Caprimulgus carolinensis	Northern Cardinal	Cardinalis cardinalis
Red-headed Woodpecker	Melanerpes erythrocephalus	Red-winged Blackbird	Agelaius phoeniceus
Red-bellied Woodpecker	Melanerpes carolinus	Eastern Meadowlark	Sturnella magna
Downy Woodpecker	Picoides pubescens	Common Grackle	Quiscalus quiscula
* = non-native species		Boat-tailed Grackle	Quiscalus major

2.5 Listed Species

Official lists of rare and endangered species are produced at the federal level by the USFWS and the National Marine Fisheries Service (NMFS) and at the State level by the Florida Fish and Wildlife Conservation Commission (FWC) and the Florida Department of Agriculture and Consumer Services (FDACS). FNAI produces a list of rare and endangered species, and maintains a database of occurrences of these species in Florida. The Institute for Regional Conservation (IRC) also ranks native plant species by conservation status in the 10-county area of South Florida. The following subsections (2.5.1 and 2.5.2) discuss the listed, rare and protected plant and animal species found within the Pepper Ranch Preserve in detail.

2.5.1 Listed Plant Species

The Florida State Statute titled "Preservation of Native Flora of Florida" (Statute 581.185) provides the following definitions:

- Endangered plants mean species of plants native to the state that are in imminent danger of extinction within the state, the survival of which is unlikely if the causes of a decline in the number of plants continue, and includes all species determined to be endangered or threatened pursuant to the federal Endangered Species Act of 1973, as amended, Pub. L. No. 93-205 (87 Stat. 884).
- Threatened plants mean species native to the state that are in rapid decline in the number of plants within the state, but which have not so decreased in such number as to cause them to be endangered.
- Commercially exploited plants mean species native to the state, which are subject to being removed in significant numbers from native habitats in the state and sold or transported for sale.

There are fourteen (14) plant species at Pepper Ranch Preserve that are listed by the FDACS, three (3) as endangered, eight (8) as threatened, and 3 as commercially exploited (Table 7). One species, *Tillandsia x smalliana* is listed in Table 7 because it is a hybrid between two endangered species but is not itself listed by the FDACS. A brief description of the species listed in Table 7 and their status is included in the following paragraphs.

Two (2) additional plant species found at Pepper Ranch Preserve are designated as critically imperiled in South Florida (SF1) by the Institute for Regional Conservation (IRC). IRC is a not-for-profit organization dedicated to the protection, restoration, and

long-term management of biodiversity on a regional basis, and to the prevention of regional extinctions of rare plants, animals, and natural communities. This designation refers to the extreme rarity (five or fewer occurrences, or fewer than 1,000 individuals) of a species, or its extreme vulnerability to extinction due to some natural or human factor.

Table 7: Listed Plant Species Detected at the Pepper Ranch Preserve			
Common Name	Scientific Name	Status	
Cardinal airplant	Tillandsia fasciculata var. densispica	Е	
Giant airplant	Tillandsia utriculata	Е	
Meadow jointvetch	Aeschynomene pratensis	E	
Catesby's Lily	Lilium catesbaei	T	
Everglades palm	Acoelorraphe wrightii	T	
Leatherleaf airplant	Tillandsia variabilis	T	
Long strap fern	Campyloneurum phyllitidis	T	
Needleroot airplant orchid	Harrisella porrecta	T	
Northern needleleaf	Tillandsia balbisiana	T	
Simpson's stopper	Myrcianthes fragrans	T	
Twisted airplant	Tillandsia flexuosa	T	
Florida butterfly orchid	Encyclia tampensis	CE	
Royal fern	Osmunda regalis var. spectabilis	CE	
Cinnamon fern	Osmunda cinnamomea	CE	
Oak mistletoe	Phoradendron leucarpum	SF1	
Quillwort arrowhead	Sagittaria isoetiformis	SF1	
Reddish wild pine (native hybrid)	Tillandsia x smalliana		

E: Endangered, T: Threatened, CE: Commercially Exploited

SF1: Critically imperiled in South Florida (as designated by IRC)

Five (5) of the fourteen listed plant species found on the Pepper Ranch Preserve are classified as bromeliads. Bromeliads are members of the pineapple family (Bromeliaceae). While some of these species may be found growing terrestrially, most native bromeliads found in Florida are found growing attached to tree trunks and branches and may therefore be referred to as epiphytes (a plant that lives upon other plants; from Greek "epi" = upon "phyte" = plant). The leaves and/or roots of these airplants (depending on the species) absorb the water and nutrients they need from the air and from the rain that falls through the canopy of the tree on which they are found. Since epiphytes use their roots only to anchor themselves to another plant, they are considered non-parasitic.

Even though the 5 listed bromeliad species found on the Pepper Ranch Preserve are fairly common in the state, they are listed due to illegal collecting and the destruction of the habitats in which they are found. Additionally, infestation by the introduced Mexican bromeliad weevil (*Metamasius callizona*) has been implicated in the decline of many airplant populations around the state. Currently, there are no control measures in place for the Mexican bromeliad weevil however, close research and monitoring is taking place.

Cardinal Airplant (*Tillandsia fasciculata*), is also known as common wild pine and stiff-leaved wild pine. *T. fasciculata* is listed as an endangered plant by the State of Florida and has been recorded in 24 counties throughout Florida (Wunderlin & Hansen

2008). This epiphyte was frequently found in South Florida before the introduction of the Mexican bromeliad weevil. Today, it may be found in hammocks, cypress swamps and pinelands.

Like most of the other bromeliads in Florida, this species is often referred to as a "tank" bromeliad because the leaf axils and central stems form a "tank" or reservoir at the base of the plant. These reservoirs capture and hold water, dead and decaying plant matter (leaves, seeds, twigs, etc.), and dead and drowning non-aquatic insects; these trapped items provide nutrients for the plant (Larson et al. 2006).

Giant airplant (*Tillandsia utriculata*) also known as the giant wild pine, is the largest epiphyte and is relatively common in hammocks and swamps in South Florida. It can reach 12-30 inches in height and its flower spike may be more than six feet in height. It is also listed by the State of Florida as endangered.

Meadow jointvetch (*Aeschynomene pratensis*), is endemic to Florida, meaning it occurs nowhere else in the world. It is a State endangered species that has been recorded in only four (4) southern Florida counties (Wunderlin and Hansen 2008).

Reddish wild pine (*Tillandsia* X *smalliana*), is a hybrid orchid derived from the crossing of two State endangered native orchids *T. balbisiana* and *T. fasciculata var. densispica*; it is not itself listed by FDACS. This species has been recorded in only seven (7) southern Florida counties (Wunderlin and Hansen 2008).

Catesby's Lily (*Lilium catesbaei*) is an herb endemic to the U.S. southeastern coastal plain and is listed as a threatened species in the State of Florida. It is found nearly throughout Florida and has been recorded in 50 counties (Wunderlin and Hansen 2008). In Collier County, it has only been recorded at Wet Woods Preserve, Railhead Scrub Preserve, Big Cypress National Preserve, Collier Seminole State Park, Florida Panther National Wildlife Refuge, Picayune Strand State Forest, and Pepper Ranch Preserve. Johnson Engineering found it on the preserve on October 8, 2009 in the mesic flatwoods located in the southeastern portion of management unit 8 (see management unit map in Appendix 5).

Everglades palm (*Acoelorraphe wrightii*) is a State threatened species that has been recorded in only three (3) southern Florida counties (Wunderlin and Hansen 2008). This salt-tolerant palm is at the northern limit of its range in southern Florida. It was once common here but many plants were taken for the nursery trade (Bush and Morton 1969).

Leatherleaf airplant (*Tillandsia variabilis*) is a State threatened species that has been recorded in ten (10) southern Florida counties (Wunderlin and Hansen 2008). Like other airplants described in this plan, leatherleaf airplant is typically found in hammocks and cypress swamps.

Long strap fern (*Campyloneurum phyllitidis*) is a State threatened species that is epiphytic in hammocks and swamps and can sometimes grow on rocks or on walls in limestone sinkholes where it is reduced in size (eflora – flora of NA).

Needleroot airplant orchid (*Harrisella porrecta*) is widespread in the central and southern counties of Florida and is considered a threatened species. This airplant's flowering period is between August and November. Other common names are the jingle bell orchid or the leafless orchid (Brown 2002).

Northern needleleaf (*Tillandsia balbisiana*) also known as reflexed wild pine, is an epiphytic, "tank" bromeliad and is listed as a threatened plant by the State of Florida. Wunderlin and Hansen reported this species in 22 counties throughout Florida as of 2008 (Wunderlin and Hansen 2008). Reflexed wild pine is an occasional species in South Florida and is usually found in scrub, pinelands, strand swamps, hammocks, mangrove swamps and on shell ridges/mounds.

Simpson's stopper (*Myrcianthes fragran*) is a State threatened species found in hammocks. The red flaking bark of this tree can confuse its identification with the invasive exotic guava (*Psidium guajava*).

Twisted airplant (*Tillandsia flexuosa*), a State threatened species, is less common in Florida than the other *Tillandsia* species listed in this plan, but still frequent, especially in coastal ecosystems. It has been recorded in ten (10) southern Florida counties (Wunderlin and Hansen 2008).

Florida butterfly orchid (*Encyclia tampensis*) is locally abundant in central and southern counties of Florida; it is commercially exploited. They grow on a wide variety of trees including live oak, red maple, bald cypress (*Taxodium distichum*), pop ash and pond apple. They normally flower in June or July but may also flower at other times of the year (Brown 2002).

Royal fern (Osmunda regalis var. spectabilis) is not in danger of being extirpated in Florida because of habitat loss, habitat fragmentation or attack by an exotic, invasive pest, but because of commercial exploitation. According to Nelson (2000), the fibers from the stem of royal fern have been used as a growing medium to grow orchids as well as to make ropes and nets. Additionally, this species is believed to have medicinal benefits; other parts of the plant may have been used to treat wounds and broken bones, relieve sprains and to help alleviate coughs and diarrhea. In Florida, this species is found in hydric areas such as wet flatwoods, cypress swamps, floodplains, stream banks and bogs.

Cinnamon fern (*Osmunda cinnamomea*) is widespread in swamps, wet woods and wet meadows throughout North and South America (Cobb et al. 2005). Its status as Commercially Exploited as listed by the FDACS makes it illegal to collect it in the wild but it is commercially available for native landscaping.

Oak mistletoe (Phoradendron leucarpum)

Oak mistletoe is a parasitic evergreen subshrub and despite its name it can be found growing on other broadleaf trees such as red maple (*Acre rubrum*). This plant is not listed by the State of Florida but has been recognized by IRC as a critically imperiled species for South

Florida. This is a temperate species at the southern end of its range, and it is possible it has always been uncommon in South Florida (Gann et al. 2002).

Quillwort arrowhead (Sagittaria isoetiformis)

This plant is not listed by the State of Florida but has been recognized by IRC as a critically imperiled species for South Florida. As with oak mistletoe, this is also a temperate species at the southern end of its range, and it is possible it has always been uncommon in South Florida (Gann et al. 2002).

2.5.2 Listed Animal Species

Table 5 in section 2.4.2 indicates which of the wildlife species documented for Pepper Ranch Preserve are protected by the USFWS (2009) and FWC (2009). Listed wildlife species that have been observed at Pepper Ranch Preserve to date include: Audubon's Crested Caracara, Bald Eagle, Florida Sandhill Crane, Limpkin, Little Blue Heron, Roseate Spoonbill, Tricolored Heron, Snowy Egret, White Ibis, Wood Stork, Big Cypress Fox Squirrel, Florida Black Bear, Florida Panther, American Alligator and Gopher Tortoise. The following is a brief description of the conservation status for those species occurring at the preserve that are currently listed as threatened or endangered by the State or federal government.

Audubon's Crested Caracara (Polyborus plancus audubonii)

This State and federally threatened species nests predominately in cabbage palms where it will lay 2-3 eggs in late winter. Agricultural development for improved pastures and citrus groves, as well as indiscriminant killing has contributed to the caracara's decline in Florida. It was officially listed on the federal list of threatened species in August 1987 (Kale and Maehr 1990).

Bald Eagle (Haliaeetus leucocephalus)

On June 29, 2007, the bald eagle was officially delisted and removed from the federal Endangered Species List in the lower 48 states. However, according to the USFWS Division of Migratory Bird Management, this bird of prey will continue to be protected by the Bald and Golden Eagle Protection Act, the Lacey Act and the Migratory Bird Treaty Act.

Florida Sandhill Crane (Grus canadensis pratensis)

Sandhill cranes occur in pastures, open prairies and freshwater wetlands in peninsular Florida from the Everglades to the Okefenokee Swamp. They build large nests in thick patches of vegetation in freshwater wetlands where they will typically lay two eggs. Nesting lasts from January through June (Kale and Maehr 1990). They are listed as a threatened species in the State of Florida.

Wood Stork (Mycteria americana)

This bird species was first,—sighted on the preserve by staff in 2008, and on multiple occasions since then, is listed as endangered by both FWC and USFWS. Also known as the wood ibis or flint head, this species is one of the largest wading birds found in Florida and the only stork in the United States. The wood stork is a tactile feeder and may be

found in fresh, brackish, and saltwater ecosystems. Because of its dependence on naturally functioning hydrologic systems, the National Audubon Society refers to this wading bird as the "barometer of the Everglades". For this reason, the wood stork is an excellent environmental indicator of wetland health (Mazziotti 2002).

Big Cypress Fox Squirrel (Sciurus niger avicennia)

Also known as the mangrove fox squirrel, the FWC lists Big Cypress Fox Squirrel as threatened in Florida. This species was first observed at Pepper Ranch Preserve by Wilson Miller in 2005 during a listed species survey however, none have been observed by staff since acquisition in 2009. While the species is widespread in eastern and central North America, the subspecies is endemic to southwestern Florida – specifically in the Immokalee Rise, Big Cypress Swamp, and Devil's Garden area in Collier County. Some areas of this range have become vacated, while many other suitable areas are being altered or becoming isolated through development. The subspecies uses most types of forest occurring in its range. However, dense interiors of mixed cypress-hardwood strands seem to be avoided by Big Cypress fox squirrels due to dense populations of gray squirrels (*Sciurus carolinensis*) occupying these areas. Big Cypress fox squirrels have been reported in cypress swamp, pine flatwoods, tropical hammock, hardwood hammock, mangrove swamp, and suburban habitats including golf courses and residential areas in native vegetation. Big Cypress fox squirrel densities appear to be quite low, and on this basis the subspecies can be considered inherently rare (Humphrey & Jodice 1992).

Florida Black Bear (Ursus americanus floridanus)

The Florida black bear is a subspecies of the black bear found throughout North America. Black bears have been observed on several occasions at Pepper Ranch Preserve, on neighboring properties and on Pepper Road, by staff and neighbors. Florida black bears in south Florida are listed as threatened by the FWC. Because of its large home range and low population density the black bear is particularly vulnerable to habitat loss. Even though their population is affected by illegal killing and road kills, habitat loss is the major cause of concern (Humphrey 1992).

Florida Panther (Puma concolor coryi)

This large cat is a year-round resident of undeveloped lands in South Florida including the Pepper Ranch preserve. It is listed as endangered by both FWC and USFWS. Panthers prefer hardwood hammocks and pine forests with numerous saw palmettos for resting, raising kittens, and stalking prey. Panthers are losing their habitat in South Florida and males require a large range. Increased development and traffic are another reason why this species is listed as endangered. Telemetry data from FWC demonstrates Florida panther frequently use the adjacent CREW lands and have used the Pepper Ranch Preserve on a few occasions (current Florida panther telemetry data obtained from FWC staff by Johnson Engineering, Inc. on 09/28/09). In their third revision to the Florida Panther Recovery Plan the USFWS (2008) states that there are three priority zones identified as important for panther habitat conservation: (1) Primary Zone – lands essential to the long-term viability and persistence of the panther in the wild; (2) Secondary Zone - lands contiguous with the Primary Zone, currently used by few panthers, but which could accommodate expansion of the panther population south of the Caloosahatchee River; and (3) Dispersal Zone - the area which may facilitate future

panther expansion north of the Caloosahatchee River. The Pepper Ranch Preserve is entirely within the Primary Zone for the Florida panther. Wildlife cameras have been installed throughout the preserve and have taken hundreds of photos of panther on the preserve since 2010. In 2016, panther kittens photos were also taken on the cameras.



Panther photo taken by wildlife camera on the preserve in January 2016



Panther kitten photo taken on boardwalk in December 2016

American Alligator (Alligator mississippiensis)

The American alligator is listed as <u>threateneda Species of Special Concern (SSC)</u> by FWC and as threatened by USFWS for its similarity in appearance with the endangered American Crocodile. Alligators are seen throughout the preserve on a regular basis and they are quite numerous in Lake Trafford that borders the Preserve to the south.

Gopher Tortoise (Gopherus polyphemus)

This medium-sized, native land turtle is listed by the State as a threatened species. Gopher tortoises are typically found in dry, upland habitats including scrub, xeric oak hammock, sandhills and dry pine flatwoods. Burrows are created for protection from weather, fire, and predators; they also provide refugia for more than 300 other species of animals. Active burrows exist in the pine flatwoods communities at Pepper Ranch Preserve.

2.6 Invasive, Non-native and Problem Species

In an ecological context, an invasive species is one that is aggressive in growth and expansion of range and tends to dominate others; its establishment and dominance can cause widespread harm to an ecological system by altering a plant community's species composition, susceptibility to fire and hydrology. Non-indigenous species (i.e., non-native or exotic species) are those that have been introduced purposefully or accidentally to an area outside their normal range. The characteristics of some of these species (high rate of growth/reproduction, no natural predators, easily dispersed, able to out-compete native species) make them invasive. Some indigenous species (a species whose natural range included Florida at the time of European contact circa 1500 AD or a species that has naturally expanded or changed its range to include Florida) may also become invasive. Invasions by native and non-native species often follow an alteration to ecosystem function, disruption of the food web, large-scale fragmentation of an ecosystem and/or disturbance (e.g., clearing, fire, drought, etc.) of an area. While some native species may become invasive, the establishment and dominance of non-native

species is of particular concern. The exotic plant and animal species documented within the preserve and those that have a potential to occur within the preserve are discussed in the following sections.

2.6.1 Invasive and Problem Plant Species

FLEPPC maintains a list of exotic plants that have been documented to (1) have adverse effects on Florida's biodiversity and plant communities, (2) cause habitat loss due to infestations and (3) impact endangered species via habitat loss and alteration. To date, 82 non-indigenous or non-native plant species have been detected within Pepper Ranch Preserve (Table 8), accounting for 20% of the plant species recorded there. Of the 82 exotic species, 32 are listed by FLEPPC (23 Category I and nine Category II). FLEPPC defines Category I plants as those that alter native plant communities by displacing native species, change community structures or ecological functions, or hybridize with natives. Category II plants have increased in abundance or frequency but have not yet altered Florida plant communities to the extent shown by Category I species. These definitions do not rely on the economic severity or geographic range of the problem, but rather on the documented ecological damage caused by these plants (FLEPPC 2009).

Scientific Name	Common Names	FLEPPC Category
Abrus precatorius	ROSARY PEA; BLACKEYED SUSAN	I
Ageratum conyzoides	TROPICAL WHITEWEED	
Albizia lebbeck	WOMAN'S TONGUE	I
Alternanthera philoxeroides	ALLIGATORWEED	II
Alysicarpus ovalifolius	FALSE MONEYWORT; ALYCE CLOVER	
Alysicarpus vaginalis	WHITE MONEYWORT	
Amaranthus spinosus	SPINY AMARANTH	
Asclepias curassavica	SCARLET MILKWEED	
Blechum pyramidatum	BROWNE'S BLECHUM	II I
Casuarina glauca	GRAY SHEOAK; SUCKERING AUSTRALIAN- PINE	I
	SOUR ORANGE; GRAPEFRUIT; SWEET	
Citrus x aurantium	ORANGE	
Citrus x jambhiri	ROUGH LEMON	
Commelina diffusa	COMMON DAYFLOWER	
Crotalaria pallida var. obovata	SMOOTH RATTLEBOX	
Cuphea carthagenensis	COLOMBIAN WAXWEED	
Cynodon dactylon	BERMUDAGRASS	
Cyperus rotundus	NUTGRASS	
Desmodium triflorum	THREEFLOWER TICK-TREFOIL	
Eichhornia crassipes	COMMON WATER-HYACINTH	I
Eleusine indica	INDIAN GOOSEGRASS	
Emilia fosbergii	FLORIDA TASSELFLOWER	
Eragrostis atrovirens	THALIA LOVEGRASS	
Eragrostis ciliaris	GOPHERTAIL LOVEGRASS	
Eugenia uniflora	SURINAM CHERRY I	
Eulophia graminea	(no common name)	
Ficus microcarpa	INDIAN LAUREL I	
Hedychium coronarium	BUTTERFLY GINGER	
Hemarthria altissima	LIMPOGRASS II	
Hydrilla verticillata	WATERTHYME, HYDRILLA I	
Hymenachne amplexicaulis	TROMPETILLA, WEST INDIAN MARSH GRASS	I

Formatted: Highlight

Hyptis verticillata	JOHN CHARLES	
Imperata cylindrical	COGONGRASS	I
Indigofera hirsute	HAIRY INDIGO	
Kigelia pinnata	SAUSAGE TREE	
Lantana camara	LANTANA, SHRUBVERBENA	I
Leucaena leucocephala	WHITE LEADTREE	II
Ludwigia peruviana	PERUVIAN PRIMROSEWILLOW	I
Lygodium microphyllum	SMALL-LEAF CLIMBING FERN	I
Macroptilium lathyroides	WILD BUSHBEAN	
Mangifera indica	MANGO	
Medicago lupulina	BLACK MEDIC	
Melaleuca quinquenervia	PUNKTREE	I
Melinis repens	ROSE NATALGRASS	I
Momordica charantia	BALSAMPEAR	<u>II</u>
Murdannia nudiflora	NAKEDSTEM DEWFLOWER	
Murdannia spirata var.		
parviflora	ASIATIC DEWFLOWER	

Scientific Name	ne Common Names	
Nephrolepis multiflora	ASIAN SWORD FERN	I
Oldenlandia corymbosa	FLATTOP MILLE GRAINES	
Panicum maximum	GUINEAGRASS	II
Panicum repens	TORPEDO GRASS	I
Paspalum notatum	BAHIAGRASS	
Paspalum urvillei	VASEYGRASS	
Pennisetum polystachion	WEST INDIAN PENNISETUM; MISSIONGRASS	II
Phoenix roebellini	PYGMY DATE PALM	_
Pistia stratiotes	WATER-LETTUCE	I
Pouzolzia zeylanica	POUZOLZ'S BUSH	
Pseudelephantopus spicatus	DOG'S-TONGUE	
Pseudogynox chenopodioides	MEXICAN FLAMEVINE	
Psidium cattleianum	STRAWBERRY GUAVA	I
Psidium guajava	GUAVA	I
Pteris vittata	CHINESE LADDER BRAKE	II
Richardia grandiflora	LARGEFLOWER MEXICAN CLOVER	
Sacciolepis indica	INDIAN CUPSCALE	
Salvinia minima	WATER SPANGLES	
Schinus terebinthifolia	BRAZILIAN PEPPER	I
Senna alata	CANDLESTICK PLANT	
Senna obtusifolia	COFFEEWEED; SICKLEPOD	
Senna pendula var. glabrata	VALAMUERTO	I
Solanum diphyllum	TWOLEAF NIGHTSHADE	II
Solanum viarum	TROPICAL SODA APPLE	I
Spermacoce verticillata	SHRUBBY FALSE BUTTONWEED	
Sphagneticola trilobata	CREEPING OXEYE, WEDELIA	II
Sporobolus indicus var.		
pyramidalis	WEST INDIAN DROPSEED, SMUTGRASS	I
Syzygium cumini	JAVA PLUM	Ī
Thelypteris dentate	DOWNY MAIDEN FERN; DOWNY SHIELD FERN	
Thunbergia grandiflora	SKYVINE	
Tradescantia zebrine	WANDERING-JEW; INCHPLANT	
Trifolium repens	WHITE CLOVER	
Triumfetta semitriloba	SACRAMENTO BURRBARK	

Urena lobata	CAESARWEED	<u>I</u> H
Urochloa distachya	TROPICAL SIGNALGRASS	
Verbena brasiliensis	BRAZILIAN VERVAIN	

As of the February 2009 acquisition of the Pepper Ranch Preserve by the Conservation Collier program, the most problematic non-indigenous or exotic, invasive plant species were torpedo grass, Brazilian pepper and cogon grass. To date, exotic plant treatments have taken place on approximately 1,750 acres of the preserve, the remainder of the preserve is open wetland and pasture. —within the northern SSA 7 area, a total of approximately 50 acres. The focus was primarily on small leaf climbing fern, otherwise known as Old World climbing fern. Contractors also moved and treated the Brazilian pepper growing along the main road within the preserve that leads to the oil fields. The control/removal of invasive, exotic species is discussed in detail in section 4 of this document.

2.6.2 Invasive and Other Potential Problem Animal Species

Although Florida does not have an official exotic, invasive animal species list, at least 400 exotic fish and wildlife animal species have been reported in Florida, and approximately 125 species are established.

Two non-indigenous, invasive animal species have been documented on the preserve: the brown anole (*Anolis sagrei*), and the feral pig (*Sus scrofa*). One potentially problematic species is the coyote (*Canus latrans*). Based on the natural communities found within the preserve, proximity to residential areas and geographic location, several more species (native and non-native) have the potential to impact Pepper Ranch Preserve to varying degrees and may yet be observed on site during future visits and wildlife surveys. Brief descriptions of documented, invasive, or potentially problematic species are provided in the following paragraphs.

Brown Anole (Anolis sagrei): documented within the Pepper Ranch Preserve

Also known as the Cuban anole, the brown anole is native to Cuba, the Bahamas, and neighboring islands (Schwartz & Henderson 1991). Like other anoles from the islands, this species is a small, tropical, diurnal, arboreal, territorial, and insectivorous lizard (Campbell 2001). The brown anole was first documented in the Florida Keys in the late 1800s (Lee 1985) and has since spread throughout Florida, into Georgia and into two other southeastern states (Campbell 1996). It feeds on a wide variety of insects, amphipods, and isopods. Brown anoles also prey on other small vertebrates including the hatchlings of the native green anole (*A. carolinensiis*; Campbell 2000). Campbell (2000) showed that, in the absence of the exotic brown anoles, native green anoles occupy perches from ground to the canopy of vegetation. However, in the presence of the exotic anole, native anoles move higher in trees, occupying only the trunk and crown of trees. Dietary overlap is high between both species, but the overall affects of the brown anole on the green anole are still undetermined.

Feral pig (Sus scrofa): documented within the Pepper Ranch Preserve

Hogs were first brought to Florida in the mid 1500's to provision settlements of early explorers. Their high rate of reproduction and their ability to adapt to Florida's natural

areas has led them to populate every county in the state. Today, Florida is second only to Texas in its feral hog population (Giuliano & Tanner 2005a; 2005b). While feral pigs are able to survive in a variety of habitats, they prefer large forested areas interspersed with marshes, hammocks, ponds, and drainages; cover in the form of dense brush; and limited human disturbance (Giuliano & Tanner 2005b). Dense cover is used as bedding areas and provides protection from predators and hunters. Feral pigs are omnivorous, opportunistic feeders consuming grasses, forbs, and woody plant stems, roots, tubers, leaves, seeds, fruits, fungi and a variety of animals including worms, insects, crustaceans, mollusks, fish, small birds, mammals, reptiles, amphibians and carrion. Their propensity for digging for foods below the surface of the ground (rooting) destabilizes the soil surface, resulting in erosion and exotic plant establishment. Additionally, this behavior uproots or weakens native vegetation (Giuliano & Tanner 2005a; 2005b). Due to the natural communities that are found within the preserve, this species has the potential to thrive within the boundaries. As these animals are highly visible outside of natural plant communities, adjoining residents of the preserve may be useful in the early detection of this nuisance animal. Several large families of feral pigs have been observed on the preserve since its acquisition.

Coyote (Canis latrans): documented within the Pepper Ranch Preserve

Covotes may have potential to become a problematic species. Covotes were introduced in very small numbers to Florida during the 1920's for sport hunting with domestic dogs. This introduction did not lead to the establishment of coyote populations in Florida. Concurrently, these canids expanded their range eastward across the United States and Canada as a result of nonspecific needs in habitat and food, decreased competition from other predators, large litter sizes and anthropogenic changes to the landscape. Since many species naturally expand or change their home ranges in response to climate and resource availability. The coyote may be considered naturalized or native to Florida based on fossil records and natural range expansion (FWC 2007). This crepuscular (active mostly at dawn and dusk) species is elusive and may travel individually or in groups of two or three (Coates et al. 1998). Coyotes commonly enlarge burrows made by other animals such as armadillos or gopher tortoises to use as dens or use dense vegetation for cover. Coyotes will kill smaller predators (e.g., foxes, opossums, etc.) and will help control rodents, which can be beneficial to turkey, quail, ducks and other ground nesting birds. Because their food habits are diverse, coyotes are unlikely to significantly affect the population of any single species. Coyotes are not reported to interact with bobcats and do not appear to influence bobcat home range size. Coyotes may prove beneficial in controlling potential problem species such as feral cats and hogs (Thornton, Sunquist, and Main 2004). Coyotes may attack calves (FWC 1999-2010), however this is relatively uncommon in Florida.

3.0 Previous Conditions of the Preserve; Current Use of the Preserve and Adjacent Land Uses

3.1 Previous Land Uses of the Preserve

The earliest aerial photographs obtained of the preserve were taken in 1940, 1953 and 1963 (Figures 8, 9, and 10). Digital images were downloaded from the U.S. Department of Interior USGS historic aerial photo web page (USDI 2004) and the Florida Department of State Aerial Photography of Florida web page (FDOS 2006) and georeferenced in ArcGIS 9, ArcMap Version 9.3. Aerial photographs (1975 – 2009) from the Collier County Property Appraiser web page were also reviewed.




Figure 8: 1940 Aerial View of Pepper Ranch Preserve

Review of the historic aerial photographs revealed Lake Trafford Road and Pepper Road (both inside and along the edge of the preserve) existed in 1940. In 1940, the majority of the preserve was natural with the exception of one agricultural field in the southeast portion of the preserve in Section 26, Township 46 South, Range 28 East (identified as Folio Nos. 00052680009 and 00052640007 on the Collier County Property Appraiser web site) and one agricultural field in the northernmost central extent of the preserve in Section 22, Township 46 South, Range 28 East (identified as Folio No. 00052360002 on the Collier County Property Appraiser web site). Two areas cleared for homesteads appear on the 1940 aerial photograph in the southeastern extent of the preserve, one of

which is the area surrounding the current caretaker house located at the main preserve entrance on the south side of Pepper Road. A trail from the homestead to Lake Trafford is evident on the 1940 aerial photograph. The other homestead appears on the 1940 aerial photograph east of the current caretaker house near Pepper Road along the northeast boundary of Section 35.




Figure 9: 1953 Aerial View of Pepper Ranch Preserve

Between 1940 and 1953, no further development was evident in the preserve. Between 1953 and 1963, the majority of the site agricultural fields and associated ditches were created. Between 1963 and 1975, a crescent shaped agricultural field was added at the central western extent of the site agricultural fields. Between 1975 and 1980, the eastwest portion of Trafford Oaks Road was constructed, dividing the marsh and forested wetlands through which it was constructed.




Figure 10: 1963 Aerial View of Pepper Ranch Preserve

Prior to 1995, the only additional development in the preserve included the construction of three oil fields adjacent to the hydric improved pasture at the central western extent of the preserve and the construction of two agricultural fields in the most eastern extent of the preserve (in the northeast corner of Section 35). Two of the three oil fields are currently active. The two agricultural fields have been fallow since approximately 2006 and are currently overgrown with shrubs (i.e. Brazilian pepper, wax myrtle, etc.).

3.2 Previous Land Uses of Adjoining Properties

Based on review of the 1940 aerial photograph, the lands that adjoin the preserve were natural. In 1953, canals were excavated from Lake Trafford and agricultural fields were constructed around them. Natural plant communities were converted to agricultural fields on lands southeast of Lake Trafford and south of Lake Trafford Road east of the preserve. Adjoining lands west and southwest of the preserve, the majority of which are now part of the major wetland slough on CREW lands, remained undeveloped.

By 1963, agricultural fields and residential properties (± 2.5 or ± 5.0 acre properties north and south of Lake Trafford Road) were constructed to the east between the preserve and the town of Immokalee. By 1975, a large agricultural field was constructed northeast of the preserve. The wetland slough (currently on CREW lands) and natural communities immediately north and south of the preserve remained undeveloped. Between 1980 and 1985 Trafford Oaks Road was extended from its western terminus to the south. Estate-

sized residential properties along the south extension of Trafford Oaks Road were developed with homes and borrow ponds. These properties adjoin the southwestern extent (Section 33) of the preserve. By 1985, agricultural development surrounding Lake Trafford had increased, however, the major wetland slough to the west and northwest of the Lake and the wetland slough south of the Lake remained undeveloped. Throughout the 1990s and early 2000s, residential development continued along Lake Trafford Road east of the preserve and agricultural uses remained on lands northeast of the preserve.

In 1990, SFWMD purchased the lands that encompass the major wetland slough located west and northwest of the preserve. These lands are referred to as the Corkscrew Regional Ecosystem Watershed or CREW. Through its adjacency to CREW project lands, the preserve is connected to several thousands of acres of preserved land in southeast Lee County and Northwest Collier County, including diverse systems located in Corkscrew Swamp Sanctuary, Panther Island Mitigation Bank, other CREW lands, the Southwest Florida Regional Airport 7,000-acre mitigation site and Conservation Collier's ±367-acre Caracara Prairie Preserve.

3.3 Current Land Uses of the Preserve

Currently, there is a cattle lease with Lake Trafford Ranch, LLLP and a mineral rights lease with Newport Oil on the Pepper Ranch Preserve. The current cattle lease started in May 2014 and encompasses 2,012.10 acres of the preserve. The lease and-is for a fivethree year period with option to renew for two additional terms of one year. It brings in revenue for the program each year. If the current cattle lease is not renewed, the County will publicize a request for proposal (RFP) to the public to ensure that the current lease is replaced, in order to ensure a fair process for bidding on cattle leases. The oil drilling lease covers the two quarter sections in which the oil wells exist (southwest quarter of Section 28 and northwest quarter of Section 33, both in Township 46 South, Range 28 East). The rights reserve all minerals below 250 feet. In addition, staff will consider inviting beekeepers by publicizing a request for proposal (RFP) as a permitted use on the Preserve.

PreviouslyTo date, Conservation Collier staff has held two public outreach events per year at the preserve. The initial outreach event was held on May 9, 2009. The guided hikes offered to the public during the initial outreach event were completely filled. The second public outreach event was held November 21, 2009. Over 300 people attended the event and participated in the guided hikes and van tours of the preserve as well as the historical presentation about Pepper Ranch. Since then staff has partnered with the Immokalee One-by-one foundation to hold an Earth Day Festival in 2016 and 2016. This event was a huge success and brought hundreds of people from the town of Immokalee to the preserve. As evident from the public outreach events, there is strong public interest in Pepper Ranch Preserve.

The preserve will be open every Friday and non-hunt or holidayen Saturdays and

Formatted: Justified

Formatted: Font: Times New Roman, 12 pt, Not Italic

<u>Sundays</u> from November through the end of <u>June-April</u>. <u>and-This schedule will continue</u> each year until adequate staffing or security is allocated. -Then, additional days <u>may will</u> be added. <u>When open, tThe</u> public will be able to utilize the lodge hiking trail, walk out to Lake Trafford, and picnic next to the lodge. The public will also be allowed to obtain a daily permit that will allow them to drive through the northern portion of the Preserve staying on the main road only until safe trail systems can be created. Public use of the preserve must be consistent with the preserve management goals and is discussed in section 4 of this document.

3.4 Current Land Uses of Adjoining Properties

The Pepper Ranch Preserve is bordered on its west and northwest boundaries by the CREW project lands. These are lands purchased by the SFWMD under the Save our Rivers program. Adjacent to the west are CREW project lands known as the CREW Marsh; to the north are SFWMD lands, agricultural lands and orange groves; to the east are SFWMD and residential lands (town of Immokalee); and to the south are Lake Trafford, estate-sized residential properties (Trafford Oaks), and agricultural and undeveloped lands owned by Baron Collier Investments, Ltd.

Directly south of CREW project lands and connected to them are private conservation lands owned by the National Audubon Society (Corkscrew Swamp), more conservation lands owned by the SFWMD (Bird Rookery Swamp) and various private mitigation lands, all together encompassing 60,000 acres, of which over 42,000 acres is currently held in conservation. The SFWMD makes certain capital improvements to its lands such as fencing, access roads/trails, and may provide basic public facilities on lands. Additionally, habitat management such as exotic plant species removal and prescribed burning may be conducted. Florida Statutes (F.S. 373.59) also require the SFWMD to develop appropriate public use.

The organization most frequently associated with CREW project lands is the CREW Land and Water Trust, Inc. (CREW TR), a nonprofit environmental education organization established in 1989 to coordinate the land acquisition, land management, and public use in the 60,000-acre CREW project area. The CREW TR does not own the land but operates in partnership with the SFWMD.

Approximately 180 acres of conservation land exists along the central eastern boundary of the Pepper Ranch Preserve. A 625-acre impoundment that serves as a dredge disposal site for nutrient-laden muck from the bottom of Lake Trafford is located east of the conservation land. The Lake Trafford hydraulic dredging restoration project is being conducted by SFWMD in cooperation with the Florida Department of Environmental Protection (FDEP) and through the cooperative efforts of various local organizations and state and federal agencies. Phase I of the restoration project, completed in 2006, removed over three million cubic yards of muck from the deeper portions of the lake. Phase II and III is projected to removed several million additional cubic yards of muck from the lakeshallow littoral zone. The project was scheduled to be completed in in November 2010. 2007, but has been delayed due to lower than expected water levels.

Eight sections of land owned by Turner Grove Citrus LTD Partnership located to the northeast of the ranch, and extending into Lee County, currently have citrus groves on them.

3.5 Cultural, Historical and Archeological Resource Protection

The Pepper Ranch Preserve is within an area of historical and archaeological probability. Before conducting any development near Lake Trafford, County staff ordered a Phase I Reconnaissance Cultural Resource Survey which was conducted in November 2010 by the Archaeological and Historical Conservancy, Inc. The survey was conducted in the area surrounding the lodge/visitor center. Prehistoric and historic archaeological sites were found and the lodge/visitor center building was deemed historical. Recovered cultural materials included artifacts and faunal bone. Prehistoric remains included three sand tempered pottery sherds. One test hole uncovered a prehistoric midden site that included a component of historic refuse. Additional historical and archaeological sites are most likely present on the property. Before conducting any additional development, the County will obtain Archaeological Surveys within the area(s) to be developed. When possible, the County will refrain from building in areas identified as potential archaeological sites. If development is unavoidable in areas identified as potential archaeological sites, the County will develop improvements under the guidance of an archaeologist. In addition, the County will notify the Division of Historical Resources immediately if further evidence is discovered to suggest any archaeological or historic resources are present in areas that were not identified in the Phase I. If such resources are identified on-site, a professional survey and assessment shall be instituted. archaeologist shall prepare a report outlining results of the assessments and issue recommendations to County staff about management of any sites discovered, per provisions of the Land Development Code Section 2.2.25. This report shall be sent to the Division of Historical Resources. The County shall cooperate fully with direction from the Division of Historical Resources on the protection and management of archaeological and historical resources. The management of these resources will comply with the provisions of Chapter 267, F.S., specifically Sections 267.061 2 (a) and (b).

The <u>lodgevisitor center</u> may be considered <u>a</u> historical structure. The <u>visitor center is undergoing the process of gaining historical designation</u>. Retaining one or both these structures and their designation as historical structures may provide benefits to Conservation Collier in terms of obtaining grant funds for restoration. Staff will work with the County's Historical/Archeological Preservation Board to identify options associated with designation of the <u>visitor centerlodge</u> as <u>a</u> historical structure and provide these options to the Conservation Collier Land Acquisition Advisory Committee (CCLAAC) for recommendation to the Collier County BCC.

3.6 Major Accomplishments since Acquisition

Collier County purchased the Pepper Ranch Preserve in February of 2009. The table below lists the accomplishments since acquisition of the property.

Table 9: Major Accomplishments During Previous Years		
Accomplishment	Year(s)	
Exotic vegetation treatment in North Stewardship Sending Area (SSA) (50 acres)	2009	
Cattle Vat Cleanup	2009	
Removal of Old Structures	2009	
Creation of a New Trail by the Lodge	2009	
Two Public Outreach Events	2009	
First Youth Hog Hunt Held	2010	
Public Hog and Small Game Hunts Began	2010	
Initial Exotic vegetation treatment and maintenance began-completed in SSA (720 acres) 200940 2011		

4.0 Future Use of the Pepper Ranch Preserve including Management Issues, Goals and Objectives

This section describes the main management issues, goals and objectives for the Pepper Ranch Preserve as well as the overall management framework. Central to the management of the Preserve is the mission of the Conservation Collier Program, and the goals and objectives set forth in this management plan.

4.1 Management Plan Framework

Each property purchased by Conservation Collier shall have its own management plan. At the time the Pepper Ranch Preserve was purchased, the Conservation Collier Ordinance required that an "Interim" Management Plan be developed within 60 days of closing. Interim plans include basic items such as removal of invasive, exotic vegetation and trash, establishing site security, developing management partnerships and planning for public access. The interim plan for this site was officially approved in September 2009. The ordinance then requires a "Final" ten-year management plan be developed within two years. Subsequently, the property management plan must then be reviewed every five years. Final management plans, however, are considered living documents and can be updated at any time. Review of all management plans start in the Lands Evaluation and Management subcommittee and must be approved by both the CCLAAC and the Collier County BCC.

4.1.1 Preserve Manager: Contact Information

The site manager for Pepper Ranch Preserve will be a designated Collier County Environmental Specialist who may be contacted through electronic mail: ConservationCollier@Colliergov.net.

4.2 Planned Uses and Assessment of their Impacts

While visitor attendance increases every year, Future planned public uses will be consistent with the primary goals of conservation, preservation, restoration and maintenance of the resource. Official public use of the entire preserve will not be possible until safe public access trails can be created. However, citizens that desire to visit the northern public use area may do so by signing a waiver or by obtaining a daily use permit on days that are open to the public. These will allow them access at their own risk and releases the liability of the County until safe access is established. Details of publiclanned uses for the Pepper Ranch Preserve and an assessment of their potential impacts are provided in the following sections.

4.2.1 Identification of Public Uses Consistent with Preservation, Enhancement, Restoration, Conservation and Maintenance of the Resources

The Conservation Collier Ordinance 2002-63 constrains the use of this property to "primary objectives of managing and preserving natural resource values and providing appropriate natural resource-based recreational & educational opportunities." Natural resource-based recreation shall mean all forms of uses, which are consistent with the goals of this program, and are compatible with the specific parcel (Ord. No. 02-63, as amended§ 5, 12-3-02). Additionally, no dumping, use of unauthorized vehicles, or removal or destruction of natural or historical/archaeological resources will be permitted within the preserve. The goal is to allow limited, non-destructive public access to native plant communities and animal species. Currently, the preserve rules are those identified in Collier County Ordinance 76-48 (available from www.municode.com), as amended.

The following are *consistent* uses for this particular site: hiking, nature photography, camping, horseback riding, bird watching and hunting. *Inconsistent* uses include off_road vehicle use (ORV), cell phone towers, shooting ranges, and the recreational use of drones.

There is one lease and a number of easements existing on the Pepper Ranch Preserve, as identified below (see Figure 11):

Lease:

A cattle lease held by Lake Trafford Ranch LLP for Two Thousand Twelve points One (2,012.1) acres of property, as described in Appendix 6, for the sole purpose of cattle grazing and incidental activities that are directly related to beef cattle production for a term of three (53) years, commencing on May 27, 2014, with two 1 year renewal options February 2, 2009, with payments, terms and provisions as set forth in Cattle Lease, attached as an Exhibit to the Pepper Ranch Purchase Agreement. This lease brings in revenue for the property management. Staff also obtained a range management study from Natural Resources Conservation Service (NRCS) to further direct cattle lease operations at Pepper Ranch Preserve. The current lease and every lease thereafter should abide by the best management practices outlined in the current NRCS range management plan for the property.

Formatted: No bullets or numbering

Formatted: Highlight

Formatted: Font: Not Italic

Formatted: Highlight

Formatted: Font: 12 pt, Font color: Black

Easements:

- Access Easement entered into on February 6, 2009, with Lake Trafford Ranch LLP for a 30' wide strip of land running along the main interior ranch road, following an overall east to west directional track, and leading from the main ranch gate to the oil wells situated along the western side of the ranch. Recorded in O.R. Book 4425 and Page 3302, Public records of Collier County. The grantee is responsible for maintenance of this easement.
- Access Easement entered into February 2, 2009 by Lake Trafford Ranch LLP in favor of Baron Collier Investments (BCI), Ltd., a Florida Limited partnership, over a 15' wide strip of land running over the same main interior access road as the above easement but before arriving at the oil wells, turning south to facilitate access to a parcel adjoining the southern boundary of the Pepper Ranch Preserve. Access is granted solely for purposes of ingress and egress to serve specific activities on the BCI lands, which are cattle grazing, ranching, hunting and forestry. Recorded in O.R. Book 4425, Page 3263, Public Records of Collier County. The grantee is responsible for maintenance of this easement.
- Stewardship Easement Agreement recorded in OR Book 4089, Page 3837, Public Records of Collier County.
- Drainage Easement for 40' along SE corner of property in Section 35, recorded in O.R. Book 49, Page 147, Public Records of Collier County.
- Access Easement in favor of Trafford Oaks for 60' as for portion of Trafford Lakes Road that traverses Pepper Ranch property, as recorded in O.R. Book 907, Page 1383, Public Records of Collier County.
- A Conservation Easement over portions of the property associated with panther and/or wetland mitigation will be granted to the South Florida Water Management District (SFWMD).

Conservation Collier staff will maintain an open line of communication with the oil well operators to develop operational protocol where needed and to ensure this operation continues in a safe and clean manner at the preserve. This level of coordination will also be extended to the cattle lease holder on land management activities at the preserve. Staff will also obtain a range management study from NRCS to further direct cattle lease operations at Pepper Ranch Preserve.

An apiary lease is being considered for the preserve. This will be advertised for bidding to the general public and will be approved by the BCC before implementation. A minimal amount of hives will be allowed to be placed on the property away from public use areas.

No other easements, concessions or leases exist on Pepper Ranch Preserve or are proposed for the future, unless they further conservation objectives, such as a conservation easement.




Figure 11. Pepper Ranch Preserve Easement and Overlay Map

4.3 — Desired Current and Future Desired Conditions

This section includes a description of the <u>current and</u> proposed future conditions for the site's natural areas. Management techniques to achieve these conditions are outlined in section 4.4.

After managers complete recommended management actions, Pepper Ranch Preserve will consist of upland mixed forest, strand swamp, slough, prairie hammock, pine flatwoods, dry prairie, freshwater marshes, bottomland forest, and wet prairie. These communities will have a similar structure and composition to those that existed before non-indigenous people settled the region and before the exclusion of fire. Through restoration efforts the site will be vegetated with appropriate native flora that will provide suitable cover for a variety of wildlife species.

4.4 Goals for the 10-year period 2010-2020

A set of goals and objectives for Pepper Ranch Preserve were developed in conjunction with the drafting of this Management Plan. The goals and objectives in this plan are tailored specifically for Pepper Ranch Preserve based on the purposes for which the lands were acquired, the condition of the resources present, and the management issues for the property. On-site managers should be familiar with this entire Management Plan. Goals

and objectives from the Interim Management Plan for the Pepper Ranch Preserve were reviewed to determine whether they should be included in this plan. The goals and objectives presented here reflect programmatic goals and ideas of Conservation Collier personnel in charge of managing and protecting the area. These goals shall not be modified, but specific application of management techniques may take into consideration input by user groups and other stakeholders from outside the program, accommodating user needs and desires where practicable and where overarching management goals are not violated.

Management issues are discussed below in separate sections. Within each section, approaches for dealing with these issues are described. The ability to implement the specific goals and objectives identified in this plan is dependent upon the availability of staffing and funding sources. The following goals have been identified for Pepper Ranch Preserve:

- **Goal 1:** Eliminate or significantly reduce human impacts to indigenous flora and fauna
- **Goal 2:** Develop a baseline monitoring report
- **Goal 3:** Remove or control populations of invasive, exotic or problematic flora and fauna to restore and maintain natural habitats
- Goal 4: Create a Prescribed Fire Plan
- Goal 5: Restore native vegetation
- Goal 6: Develop and monitor a plan for public use
- Goal 7: Facilitate uses of the site for educational purposes
- Goal 8: Provide a plan for security and disaster preparedness
- **Goal 9:** Provide preliminary Panther Habitat Unit (PHU) calculations and a draft Monitoring Plan per USFWS requirements for an onsite Panther Conservation Bank

GOAL 1: ELIMINATE OR SIGNIFICANTLY REDUCE HUMAN IMPACTS TO INDIGENOUS FLORA AND FAUNA

Action Item 1.1 Maintain the existing boundary fence and access gates on the Pepper Ranch Preserve as needed.

Currently, a fence is present along most of the Pepper Ranch Preserve boundary with the exception of along the western boundary which this preserve shares with the adjacent CREW lands, also there is no fence along the southeastern boundary along theon Lake Trafford shoreline. Under the existing cattle lease the lessee is responsible for the installation and maintenance of all fences on the preserve necessary for retaining cattle on the property. Firebreaks will be installed along fence lines that exist along upland areas. This will also allow for better access for fence line patrolling and maintenance.

<u>Action Item 1.2</u> Install signs encouraging people to stay on public access trails situated on the preserve.

Signs will be posted along public access trails to remind visitors to remain on the trails for their safety and the protection of the natural resources of the preserve.

Action Item 1.3 Identify locations of rare and listed native plant species.

The location of these species has been identified using a global positioning system (GPS) device and mapped to allow staff to monitor them. All future sightings of such plants will be GPS-located and mapped accordingly. Public trails will be constructed to avoid areas where rare and listed species exist. These locations will not be shared with the public to protect these rare plants.

<u>Action Item 1.4</u> Enforce regulations prohibiting trash or dumping- in or near the preserve.

Staff will monitor the preserve on a regular basis and if dumping occurs, enforcement actions will be sought through the County Sheriff's Department.

Action Item 1.5 Identify actual and potential locations of resident animal life and take steps such as locating visitor amenities away from animal nesting sites.

An inventory of sensitive areas, such as location of listed plant species and animal nesting sites, will be maintained based on existing knowledge and to be built upon with all future protected species surveys that are conducted at the preserve. During the development of public use facilities this inventory will be utilized to locate the amenities away from known sensitive areas.

<u>Action Item 1.6</u> Avoid non-target damage to native plants and animals, especially rare species, during invasive, exotic plant treatments.

If the use of herbicides is appropriate during the treatment of invasive, exotic plant species, decisions on the types of herbicides utilized will be made on the best information available at the time of exotic removal. Staff has prohibited the use of herbicides containing Imazapyr (e.g., Arsenal) due to reports that these herbicides have potentially caused a great deal of non-target damage throughout the state. Licensed County or State contractors will be monitored closely to ensure the proper herbicide applications are being utilized while treating the site. In addition, close attention will be taken to identify listed species (Table 7) that may be attached to invasive trees being cut down or removed. Individuals of these species will be relocated prior to removal. Special attention will be given to avoid damage to native species in the vicinity of exotic removal activities.

Action Item 1.7 Note, research and provide input as to all site development occurring adjacent to Pepper Ranch Preserve to determine that the proper site development permits have been obtained and that the site development complies with the permits.

Activities on adjacent and neighboring properties may have an impact on the indigenous plant and animal life on the Pepper Ranch Preserve. As such, all existing local, state, and federal regulations should be strictly followed and enforced during any site development adjacent to the preserve. It shall be the responsibility of the developer to establish erosion control measures and vegetation protection measures (i.e., protective fencing or barriers). If any site developer working in areas adjacent to the preserve does not take the necessary control measures, construction shall be

immediately halted until control measures are put into place and mitigation and/or remediation will be the sole responsibility of the developer.

GOAL 2: DEVELOP A BASELINE MONITORING PROGRAMREPORT

Action Item 2.1 Establish a long-term biological monitoring program and conduct additional wildlife surveys.

Long-term management of the preserve should be based on biological data. Changes following baseline conditions should be assessed as negative or positive, and management strategies changed appropriately. This section discusses information needs and long-term monitoring needs.

Pro Native Consulting has conducted a floristic inventory of the Pepper Ranch Preserve; these findings will comprise the baseline floristic data on which future actions will be based. The site should be inspected by Conservation Collier staff at least twice a year and thoroughly inventoried at regular intervals (ca. 5-10 years) to detect new invasions (by natives or exotics) and extirpations. Areas undergoing extreme restoration should be assessed more frequently. While some wildlife data has been collected, additional baseline data should be collected, especially on invertebrates, small mammals, reptiles, and amphibians. The site manager may contract this work out or enlist the assistance of local educators to coordinate student research projects. Wildlife surveys, like plant surveys, should take place at regular intervals (ca. 5-10 years) to detect long-term trends.

White-tailed deer and turkey surveys will be conducted each year to aid in the County's panther mitigation requirements as well as providing staff with the number of surplus animals that will be allowed to be taken in the Quality Wildlife Management Hunt Program.

Photo points will be established throughout the preserve. The total number of photo stations installed will be dependent on future restoration plans and staffing levels. Locations of photo points will be recorded with a GPS unit and all photographs taken at these locations will be taken at a standard height and angle of view. During photo documentations, one photo will be taken in each of the cardinal directions (north, east, south and west) and a 360-degree panoramic photo will also be taken. Photos will be taken with a vegetation profile board to aid in the determination of what (if any) changes occur over time. These photos will help to monitor exotic removal efforts and native plant recruitment, as well as the result of other land management activities. If necessary, more photo points will be established to aid in management decisions.

<u>GOAL 3:</u> REMOVE OR CONTROL POPULATIONS OF INVASIVE, EXOTIC OR PROBLEMATIC FLORA AND FAUNA TO RESTORE AND MAINTAIN NATURAL HABITATS

Action Item 3.1 Prioritize the removal of invasive, exotic and/or problematic plant species.

Due to the size of Pepper Ranch Preserve, it will be helpful to the preserve manager to prioritize the exotic control efforts by area of the preserve. The preserve has been divided up into 3 management phases 1-3. A management unit map has been created (Appendix 5) dividing the preserve into 9 separate management units. In general, the management units will assist the preserve manager in prioritizing and allocating resources available for the management of Pepper Ranch Preserve.

<u>Action Item 3.2</u> Acquire services of licensed and qualified contractor(s) for the removal of invasive, exotic and/or problematic plant species.

The following table (Table 10) describes recommended controls (Langeland & Stocker 2001; Langeland 2008) of the Category I, invasive, exotic plant species recorded to date on the Pepper Ranch Preserve. These recommended control methods may be altered by site managers dependent on new information and products available on the control of these species.

Table 10: Invasive, Exotic Plant Species Control Plan for the		
Pepper Ranch Preserve FLEPPC Category I species ¹		
Common Name	Recommended Control(s) ²	
Rosary pea; blackeyed	Treat base of vine with 10% Garlon 4. Site must be revisited several times to pull seedlings.	
susan	Basal bark treatment with 10% Garlon 4. Cut stump treatments are also effective with 50%	
woman's tongue	Garlon 3A or 10% Garlon 4. Small seedlings can be hand-pulled.	
	Basal bark treatment with 10% Garlon 4 is very effective, as is a cut-stump treatment with 50%	
	Garlon 3A or 10% Garlon 4. When basal bark treatment is used on trees greater than 1'	
	diameter it may be necessary to slough off loose bark in the application area to prevent the bark	
	from trapping the herbicide. Addition of 3% Stalker will increase consistency on older trees.	
gray sheoak; suckering	Broadcut of 4-6 lb Velpar ULW may be used when appropriate.	
	Common Name Rosary pea; blackeyed susan woman's tongue	

Table 10: Invasive, Exotic Plant Species Control Plan for the
Pepper Ranch Preserve FLEPPC Category I species ¹ (continued)

1 epper Ranch Preserve Philip Category 1 species (continued)		
Common Name	Common Name	Common Name
		For seedlings and small plants up to ½ inch
		diameter, use a basal bark treatment with 10%
		Garlon 4. This species takes a long time to die,
		and may require a subsequent herbicide
		application. For larger stems, use a cut-stump treatment with either 50% Garlon 3A or 10%
Eugenia uniflora	Surinam cherry	Garlon 4. Seedlings should be hand pulled.
Еидениа иніјіога	Surmain cherry	Basal bark application of 10% Garlon 4 is
Ficus microcarpa	Indian laurel	effective.
Hymenachne	maian iaurei	Foliar treatment with 3-5% Rodeo.
amplexicaulis	trompetilla	Total treatment with 5 5% Rodes.
compressive courts	Hompouna .	3-4 qt. Roundup Pro or 0.5 qt. Fusulade per acre.
		For high volume, spot treatment use 3%-5%
		Roundup Pro. Herbicides should be used in
		combination with burning or tillage for optimum
		control. See IFAS publication SS-AGR-52 for
Imerata cylindrical	cogongrass	additional information.
		Treatments can be basal bark, foliar and/or cut
		stump, depending on the size of the plant, with
		Renovate 3 in aquatic conditions or Garlon 4 in
7 7	D	upland areas. Adjust percentage of chemical
Ludwigia peruviana	Peruvian primrosewillow	based on application method. Thoroughly spray foliage to wet with 1.25%
		Garlon 4 (4 pt/acre), 0.6% Roundup Pro
		(maximum 5 pt/acre), 1.0%-3.0% Rodeo
		(maximum 7 pt/acre). Only Rodeo can be used if
		plants are growing in aquatic site. Plants
		growing high into trees, cut vines and treat
Lygodium		lower portions. Do not apply when plants are
microphyllum	small-leaf climbing fern	under environmental stress.
		For seedlings and saplings: (1) hand pull, being
		sure not to break plant off of root system and
		remove or place in piles to help reduce the
		chance that they will re-root or; (2) Treat with
		foliar, low volume spot application of 5%
		Rodeo. For mature trees: (1) Fell large trees with
		chain saw leaving a level surface, or fell small
		trees with machete and treat with triclopyr or
		glyphosate products according to frill and girdle directions on SLN. Use aquatic versions where
		standing water is present. Monitor for
		resprouting and retreat as necessary. (3) Mature
Melaleuca	Melaleuca, punktree, paper	trees are very difficult to control with foliar
quinquenervia	bark	applications.
quinquenervia	Ualk	applications.

Table 10	: Invasive, Exotic Plant	t Species Control Plan for the
Pepper Ranch Preserve FLEPPC Category I species ¹ (continued)		

Scientific Name	Common Name	Recommended Control(s) ²
		Foliar application of 1-2% Roundup will
		provide control. Roundup (glyphosate) is a
		short-term solution, because regrowth from
Melinis repens	rose natalgrass	seed is rapid. ³
		A foliar application of Roundup at 1.5%
		provides control. Follow-up applications are
Nephrolepis brownii	Asian sword fern	necessary.4
		Foliar application of 0.75%-1.5% Rodeo
		and surfactant solution. Re-apply as
		necessary when plants re-grow to within 4-6
		inches in height; or foliar application of
Panicum repens	torpedo grass	0.5% spot treatment.
		Foliar application with endothall, diquat, or
Pistia stratiotes	water-lettuce	rodeo
Psidium cattleianum	strawberry guava	Basal bark application of 10% Garlon 4.
Psidium guajava	Guava	Basal bark application of 10% Garlon 4.
		Cut-stump treatment with 50% Garlon 3A, 10%
		Garlon 4 or a basal bark application of 10%
		Garlon 4. Foliar application of Garlon 4, Garlon
		3A, Roundup Pro, Roundup Super Concentrate,
		or Rodeo, according label directions may be
		used where appropriate. Glyphosate products are less effective when used alone in spring and
Schinus		early summer. Use Rodeo where plants are
terebinthifolia	Brazilian pepper	growing in aquatic sites.
Senna pendula var.		Foliar application, spray to wet with 1-2%
glabrata	valamuerto	Roundup Pro. ⁵
Solanum diphyllum	TWOLEAF	Foliar application of 1% Garlon 4 or 3%
	<u>NIGHTSHADE</u>	Roundup.
		Foliar application of 1% Garlon 4 or 3%
Solanum viarum	tropical soda apple	Roundup.
		Mature trees may take up to 9 months to
		die. Cut-stump treatment with 50% Garlon
		3A or 10% Garlon 4, or use a basal bark
Syzygium cumini	Java plum	treatment with 10% Garlon 4.

¹ FLEPPC 2009: Category I plants are those that alter native plant communities by displacing native species, change community structures or ecological functions, or hybridize with natives (FLEPPC 2009)

² All species except as cited otherwise

³ (Stokes 2009)

Formatted Table

⁴⁽Langeland 2008)

⁵(Langeland et al. 2003)

<u>Action Item 3.3</u> Acquire services of licensed or qualified contractor(s) for the removal of invasive, exotic or problematic animal species.

To date, three (2) introduced animal species have been documented on the Pepper Ranch Preserve, the brown anole and the feral hog. One potentially problematic species is the Coyote. No attempts will be made to remove occurrences of the brown anole at the preserve. This is a pervasive animal that is now present in natural and urban areas alike.

Widespread control of coyotes has been found to be ineffective and is not ecologically or economically defensible. Individual coyotes may need to be removed from the preserve if they become a problem to the current cattle operation; that decision will be made on a case by case basis. The coyote can be legally hunted all year long with guns, dogs, live traps, or snares. A permit is required to use steel traps, to trap on another person's property, or to use a gun and light at night. Possessing or transporting a live coyote requires a Class II captive wildlife permit and the use of poison is prohibited.

<u>Action Item 3.4</u> Implement the Quality Wildlife Management Hunt Program to assist in Feral Hog Management (See Regulations in Appendix 7-)

It is doubtful that the total eradication of this species at Pepper Ranch Preserve can be achieved, however, efforts will be made to reduce their population and limit the damage they cause to natural areas, native plants and animals. In late 2009, the Collier County BCC approved a contract with the USDA Wildlife Services for the control of feral hogs at the preserve, which was part of the property Interim Management Plan. As a result, fourteen hogs were trapped and euthanized in a short time. On January 12, 2010, Item 10D, the BCC voted to cancel the USDA contract due to public opposition and directed staff to develop a hunt program to attempt to control the hogs and to consider trapping at a later date if they cannot be controlled through normal hunting. As a result, the first Annual Youth Hunt was held at the preserve in April 2010, with the assistance of FWC, and 4 hogs were harvested as a result. The Pepper Ranch Quality Wildlife Management Hunt Program for this preserve was then developed and began on September 11, 2010. In 2010-2011, a total of two youth hunts and six public hunts were conducted. This program did provide some control of the hog population at Pepper Ranch, however only 8 total hogs were harvested during the first hunting season. Hunting alone may not properly manage the hog population and thus a monitoring program should be developed to assess the amount of hog damage to natural communities. Efforts will be made to request assistance from nearby colleges to conduct such studies. Additional control measures such as trapping may be necessary to protect the resource. Monitoring the hog population will be particularly important in the event that a wetlands mitigation bank is developed on Pepper Ranch Preserve; created/enhanced wetlands will be required to meet certain success criteria within a set timeframe and hog foraging behavior could severely impact creation/enhancement efforts. In areas where wetland restoration is to occur in the future, hog fencing may need to be installed around the restoration areas or potentially the entire perimeter of the preserve. This fencing would be paid for with mitigation funds.

GOAL 4: CREATE A PRESCRIBED FIRE MANAGEMENT PLAN

Fires were a naturally occurring event in native communities prior to mankind's intervention. The primary ecological functions of fire are to eliminate accumulated plant material, return nutrients to the soil, and germinate fire-dependent species. In today's preserve areas prescribed burning is an essential tool in both land and wildlife management, and helps reduce potential damage and hazards from wildfires in the wildland/urban interface areas. Proper prescribed burns promote the growth of green shoots, roots, and rhizomes of grasses and sedges that are then available for foraging. In wetlands, burning creates deep pools and edges for nesting and feeding of waterfowl and controls undesirable vegetation.

Much of Collier County is comprised of natural communities in general, that are dependent on fire to maintain species composition and diversity. The use of prescribed fire as a management tool will be critical to the long-term health of the natural communities and native species at the Pepper Ranch Preserve.

Action Items 4.1: Create a Prescribed Fire Management Plan

Below is the prescribed fire management plan for Pepper Ranch Preserve. The preserve land manager with assistance from the Florida Forest Service (FFS) Division of Forestry (DOF) and/or a Certified Prescribed Burn Manager will implement the prescribed fire management plan according to the specific needs of Pepper Ranch Preserve. Staff may coordinate this effort with other local qualified agencies for review and approval.

Objectives

The prescribed fire plan for the Pepper Ranch Preserve will be a program that mimics the natural fire cycle for the various natural community types identified within the preserve. Timing, based on weather conditions and ignition practices can be modified to accomplish goals ranging from exotic vegetation control to wildlife habitat enhancement and fuel reduction within burn units. This prescribed fire management plan will be implemented at Pepper Ranch Preserve for ecological purposes. The goals and objectives established for the preserve will be clearly laid out and incorporated into each prescription. Generally, prescribed burns conducted at the Pepper Ranch Preserve will involve a variety of firing techniques over a range of weather conditions to create mosaic burn patterns that will benefit an array of wildlife species.

Burn Units

The size of the Pepper Ranch Preserve, in conjunction with habitat fragmentation by existing (oil fields, cattle grazing) and future uses (lodge, possible housing, camping areas and public use trails) of the preserve create a complex mosaic of fire dependent communities. This will be taken into consideration when subdividing the preserve into burn units. The creation of burn units not only facilitates the application of prescribed fire, it will also help create a mixture of burned and unburned areas across

the preserve. Patches of unburned habitat in conjunction with newly burned areas will increase habitat heterogeneity, ensuring a wide range of habitat compositions year round for use by a diversity of wildlife species. The size and boundaries of each burn unit should be established based on the preserve boundaries and the location of existing barriers such as fence lines, ditches, roads and other existing structures. The division of burn units may change over time as the prescribed fire plan is implemented and on-the-ground logistics become more obvious. Fire breaks will consist of primitive roads, trails disked to bare mineral soil, wet lines or foam lines and/or natural vegetation breaks. When the Pepper Ranch Preserve burn plan is implemented, additional manmade barriers may be constructed as a result of the development of public use facilities.

If new fire breaks are needed, efforts will be made to minimize disturbance to existing native vegetation during their creation and maintenance, and no wetlands will be adversely impacted as a result of fire break construction. In the event of a wildfire FFSDOF may create fire breaks within existing wetlands. If plow lines are put in as a result of a wildfire, whether they are in a wetland or upland, efforts will be made to mitigate by gradinge those areas to prior grade.

Burn Frequency and Burn Season

Historically the frequency of wildfire in Florida's ecosystem varied from year to year. However, fire frequency for natural communities as found within the Pepper Ranch Preserve will generally follow these guidelines (FNAI 1990):

- wet prairies annual (1-2 year cycle) or frequent (3-7 year cycle);
- dry prairie frequent (1-4 year cycle);
- mesic pine flatwoods frequent (2-4 year cycle);
- hydric pine flatwoods frequent (3-7 year cycle);
- depression marshes more frequent around the periphery (3-7 year cycle) and becoming more occasional toward the center (8-25 year cycle);
- cypress/pine/cabbage palm transitional community from moist upland to hydric sites occasional (8-25 year cycle);
- cypress strand/dome swamp occasional around the periphery (8-25 year cycle) and rare in the deepest peat towards the center of the strand/dome (26-100 year cycle);
- slough occasional (8-25 year cycle) or rare (26-100 year cycle);
- prairie hammock occasional or rare; if oak and palm dominated on drier sites tolerate occasional light ground fires, but more diverse hammocks rarely burn;
- upland mixed forest rare or no fire; densely closed canopy limits air movement and light penetration, making high humidity relatively constant.

Burn units incorporating multiple natural communities under different fire cycles will be burned based on the community requiring the shortest cycle. The other communities within that burn unit that are on a longer fire cycle will likely not burn as frequently since fuels will not have built up. The seasonality, weather factors, or

ignition techniques of the prescribed burn will also be chosen to selectively burn the community within the unit with the shortest fire cycle.

Fire maintenance of hydric hammocks will be accomplished primarily by burning the adjacent flatwoods and marshes, reducing the fuel needed to ignite the hammock. Maintenance of natural species composition and protection from excess fuel build-up will be accomplished by allowing fire to enter the edges but not completely burn through the hammocks. Fire will be introduced into the edges of hammocks under moist conditions that will not result in a destructive fire through the hammock. Fire frequency in this situation will be dictated by the frequency of fires in adjacent communities.

Fire will be applied to freshwater marshes in conjunction with the burning of surrounding pine flatwoods to maintain open herbaceous ponds and control woody plants found primarily on the edge of these depressions. The centers of depression marshes are much wetter than the surrounding flatwoods and may not burn at the same time the flatwoods are ignited. In this case, a separate fire under guarded conditions may be needed to carry the fire across the marsh. In cypress strands, fire is beneficial for the control of hardwoods and reduction of ground fuels near their outside edge. Conditions dry enough to burn soils in the center of strands, or muck fires, would most likely be damaging to trees within them. The burning of cypress strands will take place only when moist conditions allow for light surface fires in the outer portion of the dome and avoid muck fires. Fire will be excluded from strands under dryer conditions.

Qualitative observations will be made within each burn unit on an annual basis to determine current fuel loads, habitat structure, and habitat quality. The burn schedule will then be modified as needed based on these qualitative observations. Areas where fire cannot be implemented will instead be mowed, roller chopped, or pruned to mimic effects of fire. The burn manager will conduct post-burn inspections to ensure the burn objectives are being met for each natural community. When possible, vegetation monitoring activities will be conducted around burn events to help assess the effectiveness of the prescribed burn regime.

Pile Burning

Burning of agricultural piles of vegetative debris may be conducted as needed. The piles must be placed in an open area such as a pasture and the piles must be placed at least 50 feet from a forested area or structure. A permit must be issued by the Florida Forest Service Division of Forestry. When burning restrictions are in place, the piles may only be burned by a Certified Pile Burn Manager licensed by FFS-DOF. When no restrictions are in place, the piles may be burned by trained staff, contractor or by the acting cattle manager after a permit is issued. Persons conducting the burning must have a water source large enough to extinguish the fire and a front end loader or other similar type of machine present before proceeding with burning.

Burn Schedule

Generally, prescribed burns within the Pepper Ranch Preserve will be conducted during the growing season (mid-March through early September) as well as during the dry season (November to mid-May). Essentially burns will be scheduled when conditions allow, and the timing selected to best suit the objectives for each burn unit, as well as to provide protection to listed species.

Burn Manager Duties

Florida Statute 590.125 and Chapter 5I-2 of the Florida Administrative Code (FAC) grant the FFSDOF the authority to regulate prescribed burning in Florida. Prescribed burning will be planned and carried out by a Certified Prescribed Burn Manager (as licensed by the FFSDOF) and experienced fire crews utilizing a Prescribed Burn Plan form, referred to from here on as the prescription. The planning and application of prescribed burning will comply with all applicable federal, state, and local regulations.

Each prescription will include the following at a minimum:

- purpose for the burn;
- brief description of the natural community type(s) to be burned;
- a map depicting the location of the burn, firebreak locations, potential hazard areas and escape routes for the fire crew;
- acceptable ranges of weather and soil moisture conditions;
- a pre-burn inspection of burn unit, firebreaks and any potential hazards (including power transmission lines, active cattle grazing locations, and existing manmade structures) within the burn unit;
- names and contact information for neighbors, lease holders, local fire district and other pertinent stakeholders to be contacted prior to ignition;
- techniques used to ignite the controlled burn;
- personnel, equipment and safety requirements;
- personnel assignments and responsibilities; and
- post-burn evaluation.

All necessary permits and authorizations will be obtained by the Certified Prescribed Burn Manager before implementation of the burn. As part of each prescription, the burn manager will develop an emergency action plan that will include escape routes for all personnel and actions to be taken in the event of unexpected weather changes or fire behavior.

Weather and Fuel Considerations

When developing recommendations for a prescribed burn, the burn manager will give careful consideration to weather and fuel conditions including, but not limited to: wind, relative humidity, temperature, rainfall and soil moisture, airmass stability and atmospheric dispersion. It will be the responsibility of the Burn Manager to obtain current weather forecasts from FFSDOF, and other weather sources as necessary,

prior to executing the prescribed burn. Although preferred weather and fuel conditions may vary based on specific burn objectives, Wade and Lundsford (1989) suggest the following as preferred conditions for prescribed burns in southern forests:

- 6 to 20 mph persistent surface winds;
- 30 to 55 percent relative humidity;
- temperatures above 80 degrees Fahrenheit are recommended when the primary objective is to control undesirable species;
- damp soil moistures;
- · slightly unstable or neutral airmass stability; and
- KBDI of 0 to 600 dependent on burn objectives.

Smoke Management

Smoke management is an essential component of the burn prescription. The burn manager will evaluate the potential impacts of each prescribed burn to smoke-sensitive areas located within a 20-mile radius from the location of the burn by employing a Screening System, such as recommended in Wade and Lundsford (1989). Based on definitions contained within the state regulations, smoke sensitive areas are areas within which smoke could have an adverse impact for reasons of visibility, health or human welfare (NRCS 2003). Monitoring of the prescribed burn will continue until smoke no longer presents a potential hazard and there is no potential for the fire to reignite and cause an uncontrolled fire.

Post-Burn Evaluation

The purpose of the post-burn evaluation is to ensure the objectives of the burn were attained and gain information to be used in future burns (Wade and Lundsford 1989). The post-burn evaluation will be conducted by the burn manager within one week following the burn, as well as a second evaluation after the first post-fire growing season. Quantitative vegetation monitoring, photo documentation and wildlife monitoring can be implemented to further aid in determining if the objectives of each burn were met.

Action Item 4.2 Develop Burn Units

Burn units will need to be delineated for Pepper Ranch Preserve, as outlined in the prescribed fire management plan above prior to the implementation of the plan.

Action Item 4.3 Install Perimeter Fire Lines

Fire lines will be installed utilizing best management practices to minimize impacts to mature trees, natural communities and wildlife populations. Firebreaks will be disked or mulched down to soil and will go around all mature pine trees; they will be a maximum of 8-10 feet wide.

GOAL 5: RESTORE NATIVE VEGETATION AS NEEDED

<u>Action Item 5.1</u> Evaluate the feasibility of conducting a hydrological analysis of the preserve to better determine restoration needs.

During the fieldwork conducted by Johnson Engineering, Inc. in the fall of 2009 it was noted that many of the natural wetland natural communities at Pepper Ranch Preserve are disturbed; as described in section 2.3. This disturbance appears to be hydrologic in nature due to the lack of standing water observed in these wetland communities in comparison to the undisturbed wetlands, and by the relatively high number of upland and exotic/nuisance plant species observed in the disturbed wetlands. A hydrological analysis of the preserve would provide a baseline for the development of a hydrologic restoration plan for Pepper Ranch Preserve. The presence of invasive exotic vegetation can be related to a hydrologic disturbance. Identifying hydrologic disturbances and proposing remedial measures (i.e. ditch removal or ditch blocks) at the preserve would not only work towards the restoration of natural plant communities but also possibly help limit exotic plant invasions in those areas. The analysis could be taken a step further to incorporate water quality analysis especially as it pertains to water flows into Lake Trafford.

A hydrologic analysis of the preserve would generally involve the placement of surface and ground water level monitoring wells at strategic locations throughout the preserve, mapping ditches and canals on the preserve, reviewing historic aerial photography of the preserve and determining the historic sheet flow patterns on site. This monitoring will be funded with mitigation funds.

Surface and ground water level monitoring wells installed for the purpose of this hydrologic analysis could be left in place for long-term, on-going monitoring at Pepper Ranch Preserve. The data collected would help monitor the health of wetland systems on site over time, as well as provide a baseline of wetland function that could help evaluate possible effects from proposed adjacent land use changes (i.e. if a mine was ever proposed adjacent to the preserve).

Action Item 5.2 Maintain a revised GIS map and description of FNAI natural communities and disturbed areas on the property.

Maintaining updated maps will help to guide restoration efforts.

Action Item 5.3 Plant native plant species in their appropriate habitats.

Periods following exotic removal and prescribed fire (or mechanical treatment) are essential to the recruitment of native plants. If native plant recruitment is not sufficient from the surrounding, intact seed source, efforts will be made to plant indigenous flora in appropriate habitats. Natural area restoration of Pepper Ranch Preserve should include only site-specific native plant material that has been determined to be non-problematic at the site and whenever possible, site-specific seed sources should be utilized. In addition, hardwoods that may invade the natural areas (mesic pine flatwood area) should not be planted.

GOAL 6: MONITOR DEVELOP A PLAN FOR PUBLIC USE

Action Item 6.1 Develop access and required facilities for intended public uses.

There are many opportunities for public use at the Pepper Ranch Preserve due to the size of the preserve, its proximity to the community of Immokalee and the diversity of natural communities present. In addition to general public uses at the preserve there are also revenue-generating uses as presented above, cattle lease and oil fields, hunting, apiary lease, —as well as mitigation uses that are already in place or are plannedproposed for the preserve, such as a panther conservation bank, —and Wwetland mitigation was considered but was denied by the SFWMD and Army Corp. of Engineers.—. All of these different uses considered and requested by the public may not be compatible with one another and thus a compatibility matrix was devised to better illustrate when and where at Pepper Ranch Preserve the different uses can occur. This matrix is included in this plan as Appendix 8.

Until additional staffing and funding are available, the portions of Pepper Ranch Preserve will only be open to the public on Friday, and non-hunt or holiday Saturdays and Sundays from November through June April of each year. Daily Use Permits will be required by all visitors and will be issued before access can be granted. County staff will be stationed in the visitor's center or guard house on Pepper Rd. to provide access and assistance to visitors. A temporary gate code for the electric gate will be given to visitors to access the northern access area-when they check in with the Park Ranger. A temporary combination lock will be added to the main gate to allow entry to the northern public access area until an electronic gate and/or small guard house is installed. Visitors must also check out before they leave for the day. This will ensure that all visitors are accounted for at the end of each day.—Security cameras are also facing each preserve entrance to help monitor ingress and egress. The small guard house will also be used at the north gate as a check station on designated hunt weekends.

Portable toilet facilities will be available until a permanent facility can be built. Upon professional inspection of the existing septic system associated with the lodge, it was determined to be unsafe and was crushed and permanently shut down. A new bathroom facility with showers was built in 2015 along with two—A new septic systems and a water treatment system. wells will be necessary for the development of permanent public use facilities at Pepper Ranch Preserve.

TConservation Collier staff will be discussing with the Collier County Parks and Recreation Program has been the possibility of their involvement in administering and staffing the visitor center, campgrounds and pole barn rentals since 2014? public access programs on the preserve. Staff also will also attempt to developed a volunteer program for the for all of the Conservation Collier Preserves with several master naturalists who have been to assisting with guided public tours and other activities.

To avoid impacts to natural communities at Pepper Ranch, guidelines were will need to be developed for the allowable uses on all proposed trails and other amenities.

Formatted: Highlight

Guidelines—will include instructions for users such as staying on trails to avoid altering the natural communities, and to take only pictures and leave only footprints. The__development of all trail systems at Pepper Ranch_should to the extent practicable, utilize existing trails and other impacted areas, and werebe developed along the natural edge of natural communities where their construction had/will-minimized disturbances, as well as avoided impacts to marshes and other wetland systems. The majority of the trails are not ADA accessible; however, all new trails will be evaluated for vehicle class use for ADA access.

The implementation of the proposed public uses at Pepper Ranch Preserve remains is dependent on funding, safety issues, site security and the availability of staff. The conceptual site plan (Figure 12) incorporates the following proposed components:

The South Public Access Area will be situated on the south side of Pepper Road with a parking area and trailhead that will lead to the lodge/visitors center.

- The_lodge/visitor center Conservation Collier staff has is planning to renovated the existing visitor center lodge and will use it for public use as a visitor center. The lodge could also be rented for special events. Policies will need to be created though the Ordinance, Policy and Rules subcommittee in regard to special events, lodge rentals and ecotourism. Approved eco-tours need to be compatible with this management plan.
- The lodge hiking trail The lodge hiking trail is accessible from the south public entrance. It is made up of three loops that total approximately 0.9 miles, heading east from the trailhead/parking area, meandering through oak hammock, mesic flatwoods and dry prairie communities; there iswill be a short segment of boardwalk crossing over a depressional marsh. Benches and interpretive signage existwill be placed at strategic locations along the trail.
- The boardwalk The boardwalk (length = approximately 812' or less)was constructed is proposed at the south end of the south public access area trailhead/parking area and would leads to a covered lake overlook platform. This is low priority and will only be constructed if the budget allows.
- The lake overlook platform —A covered lake overlook platform was constructed at is proposed at the terminus of the boardwalk and would allows visitors to view Lake Trafford from a raised elevation. This was built proposed to be constructed on the existing raised shoreline and not directly over Lake Trafford. A local Boy Scout added a large bench to the overlook as part of an Eagle Scout Project in 2017. EThis is also a low priority and will only be constructed if the budget allows.
- <u>CFuture eamping areas</u>- A small camping area with 10 campsites was will be developed in the current eastern pasture area located between the entrance to the south public access area and the lodge. This is will be open to the public on Friday and Saturday nights when the preserve is open, as soon as adequate staffing and/or security is available and when the bathroom facilities are completed. This campground is will be accessible to tent campers with vehicles, but not RV's. An RV may only be allowed in this campground for use by a campground host or possibly under other special circumstances.

There <u>is will be</u> no water or electricity located at the individual campsites. Camping <u>is will be</u> limited to hunters only during hunt weekends.

Officer's Trailer home or Campground host. A single-wide manufactured trailer home or an RV pad maywill be placed on a small improved area just south and west of the gate to the south public access area. It would be beneficial to have a A County Sheriff's Department officer or FWCC officer will reside there to keep watch over the property. We may also install a concrete pad with full hook-up here and will allow a campground host to live here with an RV during the months that the preserve is open to the public.

Formatted: Highlight
Formatted: Highlight
Formatted: Highlight
Formatted: Highlight




Figure 12: Original Conceptual Site Plan for Pepper Ranch Preserve




Figure 13: Current Trails as of August 2018

Formatted: Right: 0.81", Top: 1.5", Bottom: 1", Width: 11", Height: 17"

Formatted: Normal

Formatted: Font: 12 pt

Formatted: Font: 14 pt
Formatted: Left

Formatted: Font: 14 pt, Bold

The North Public Access Area is accessible from the north gate off of Pepper Road and will-provides public access for a scenic drive, hiking trails, multi-use trails, mountain bike trails and primitive camping areas. This northern area iswill be accessible after checking in at obtaining a daily use permit from the lodge/visitor center. All trails that double as firebreaks will be maintained on a regular basis, new trail creation and maintenance may be on dependent on the demand for use and available resources.

- The Scenic Drive- the public iswill be allowed to drive through the Preserve along the main access road after obtaining a free daily use permit and a temporary access code from the visitor center. This will-allows the public to view the majority of the preserve by vehicle and to view the wildlife and different ecosystems present. The driving tour is approximately 6.4 miles round trip and does not include the easement road that leads to the south property boundary. Visitors are would also be required to check out at the visitor center before they depart. During wet conditions, Until the designated trails listed below are completed, the public will be asked to keep vehiclesstay on the main roads during their tour and to drive at slow speed for safety. During normal dry conditions, When designated trails are officially open to the public, they may park in the designated trailhead parking areas.
- Hiking trails Hiking trails will provide a view of live oak hammocks, mesic flatwoods, cypress sloughs, open prairie and depressional marshes. Currently, there are five (65) differentseparate trail areas proposed totaling approximately 14.5–6.1 miles which arewill only all open to hikers hikers. However, hikers and trail runners cann also use all other trails designated as multi-use, including shared with horseback riders, or mountain bike trails with caution. Hikers must yield to bikers and horseback riders. All trails combined total approximately 23 miles not including the access roads. Many trails already exist as firebreaks. Benches and interpretive signage have will been placed at strategic locations along the trails.
- <u>SThe seasonal access hiking trail</u> <u>A This</u> 1.0-mile loop wouldill allow visitors to walk through some of the scenic wetland communities located in the western portion of the preserve during the dry season. It is located west of the scenic driving trail and will traverse through natural communities such as cypress strand and a red maple dominated bottomland forest. This trail will be limited to foot traffic to prevent damage to the sensitive wetland soils.
- Mulit-use Trails- (Hiking & Horseback riding) —Trails designated multi-use, accessible to horses and hikers, total 11.25 miles. This total includes the main access road and easement road. An area in the south central portion of the preserve will take riders though 3 miles of prairie, mesic flatwoods, oak hammock and marshes. It will also lead to the crossroads of the oil well road and the south easement road. Traveling south on this easement road will lead

Formatted: Not Highlight

Formatted: Indent: Left: 0.75", No bullets or numbering

to the southwestern most multi-use trails which are approximately 2.5 miles in length. The third trail will lead from the main road near the cattle pens north to the primitive camping area in the north central area which will total 2.2 miles round trip. The public will be required to park cars and horse trailers at the visitor center and enter through the north entrance gate or at a designated parking area at the trailheads. Equestrian use at Pepper Ranch Preserve may also require additional amenities such as watering and feeding areas. A small hand pump well may be installed in areas near riding trailheads. Riders are will be required to show documentation of a negative Coggin's test when they check in at the visitors center. Existing trails and firebreaks will be used as horse-back riding trails and in general are approximately 8-10 feet wide. The majority of the multi-use will not be shared with mountain bikers for safety reasons, however both user groups may have to pass each other on occasion on the main access roads. Signs have been will be posted to use caution when approaching horses. When horse riders are checked in at the lodge, hikers and bikers will be notified to use caution and to stay on designated trails. All multi-use trails will be maintained by the County and with help from volunteer groups.

•

Mountain biking trails- There are three main areas where mountain biking (off-road cycling) trails are existing and proposed to be created. They have been planned as Phase 1-3. The total length of the proposed trails is approximately 1315.5 miles in length, this includes the main access road, easement road and a small portion of the multi-use trails. These trails will be created in phases dependent on demand for use. The majority of the biking trails are will be very narrow in width and are kept separate from the horseback-riding trails. However, hikers and trail runners may share the mountain biking trails with caution. Two main areas will be located in the western portion of the Preserve. The first main area, Phase I, iswill be located in the west center, south of the main road that leads to the oil wells. This is will be the first completed phase of the biking trails. These trails have been named Panther Pass and Black Bear Berm. This area was formerly harvested of cabbage palms, as a result there are several existing trails were that can be used to create approximately 4.5 miles of winding singletrack trail through the forested area. _- A second area that may be created is located in the extreme southwest area, west of the easement road. This will be phase two and would be a 2 mile winding singletrack loop trail in the center of a horseback riding trail. The separate trails would not intersect and the only area where the two user groups may pass each other is on the main easement road that leads to the trails and back out. Phase 2, which has been named Kite flight three is will be aa third area would be a 5.54 mile partial perimeter trail that would run along the edges of the pastures and would start from the winding singletrack area in the west central portion of the preserve, and will run north along the pastures to the northern property boundary, then turn east and would circle through the proposed primitive camping area in the north center, then back down the multi-use trail past the cattle pens to the main road. It may then either follow

Formatted: Indent: Left: 0.75", No bullets or numbering

Formatted: Font: Not Bold

- the main road back to the parking area or will continue south of the road following the berm on the edge of the pasture all the way back to the road that leads to the parking area. Approximately, one-third of this trail has been created.
- The third possible are, may be created in the extreme southwest area, west of the easement road. This will be called phase three and would be a 2 mile winding singletrack loop trail in the center of the existing sunflower trace horseback riding trail. The bike and horse trails would not intersect and the only area where the two user groups may pass each other is on the main easement road that leads to the trails and back out. Bikers will be advised to walk their bikes past any oncoming horses on the easement road. This trail creation will be dependent on the USFWS Panther Mitigation application. They may or may not approve an additional trail in this area.
 - Trails have been and will continue to be will be created according to the International Mountain Biking Standards (IMBA) and the majority of the trails will be narrow singletrack trails that will be created and maintained by the use of hand tools such as a weed cutter and loppers etc. A local non-profit off-road cycling group called the Florida Mudcutters have been volunteering since 2012 red to assist in and have createding and maintaineding these trail systems. Helmets must be worn by bikers on these trails at all times. Trails arewill be specifically marked. Special gates may be installed in the future to allow bikers to pass through cattle gates and small bridges/crossovers may need to be built over ditches in the cattle pastures. During rainy season portions of the trails may be closed due to wet conditions. Bikers may park at the main visitor center and ride to the trails or may also park in future designated parking areas near the trailheads. The majority of the biking trails are will need to be maintained by the user group, however, County staff will determine at what level to assist based on available resources and will plan to be present on work days to supervise new trail creation.
- Primitive camping areas— PThe primitive camping may be developed after the designated pine flatwood area has had its first prescribed burn. Fuel loads are too high in this area for safe camping and campfires to be conducted. These areas will be accessible from the north public entrance. Campers may park near the cattle pens and hike up the multi-use trail to the camping area. There are three separate locations for primitive camping along the north hiking loop trail. These campsites will be walk-in only and will have no facilities.

Action Item 6.2 Pepper Ranch Quality Wildlife Management Hunt Program

The preserve has been open to the public for hunting since September 2010. Hunting has was been-limited to small game, hogs and turkey. Deer hunting wasis planned to be introduced in the Fall of 2011. The Preserve is will be closed on Saturday and Sunday during each hunt weekend. Currently, the hog hunts are limited to 10 hunters per weekend, while deer and turkey hunts are limited to 4 hunters per weekend. All hunters and hunters have designated hunting zones. The zone closest to the lake and the

Formatted: Indent: Left: 0.75", No bullets or numbering

structures is limited to archery only (See Appendix 7). <u>SeveralThree</u> successful FWC youth hunts have been held for kids age 12-17 and <u>at least</u> two are planned to be held each year depending on the continued interest and volunteers. The <u>amountnumber</u> of public hunts and youth hunts that will be allowed each year may change when other public uses of the property increase and based on wildlife management determinations.

Action Item 6.3 Recreational Drone Use is Prohibited

The use of Unmanned Aerial Vehicles (UAV), for recreational use by the general public is prohibited within the preserve. Drones have been proven to cause stampedes with horses and cattle, they can interfere with prescribed burning and wildfire operations, and they also may infringe on the privacy and safety of preserve visitors.

Requests by search and rescue organizations, fire and law enforcement agencies, other governmental and first-response agencies for a scheduled operation of non-recreational UAV on the preserve must be directed through the Division's Administrative Offices. Approval may be given for the purposes of training or reconnaissance through the Division Director.

For all other non-recreational requests related to media, land management or research, appermit may be issued on a case by case basis. Each permit application will be adequately evaluated as to the appropriateness of the requested activities and whether the use of a UAV will result in unacceptable impacts to the preserve and visitors. If a permit is issued, it will clearly identify the designated area(s) where the UAV may be operated within the park. The permit will also contain the terms and conditions to ensure safe operation and will mitigate any unacceptable impact to the resources and the public. Users will specifically be advised not to fly them in the bald eagle nesting zone west of the main campground during nesting season, or near cattle or horseback riding areas. Coordination will need to be carried out if riders are on the property. County staff can only enforce drone use when they take off and land on our property. Drones that are flown over the preserve from other properties are under the jurisdiction of the Federal Aviation Administration (FAA). All permitted users should abide by the FAA applicable laws and regulations.

GOAL 7: FACILITATE USES OF THE SITE FOR EDUCATIONAL PURPOSES

Actions Item 7.1 Develop interpretive signage to educate preserve visitors.

On <u>completedee a trail systems</u> is <u>complete</u>, site-specific signage, including directional signage, <u>has been installed will be developed</u> to educate visitors on plant identification and general ecosystem information. A <u>large sign with a map of the preserve will be installed at the north and south public entrances.</u> Additional smaller trail specific interpretive signs will be placed at the various trailheads.

Formatted: Font: Bold, Underline

Formatted: Font: Bold

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Indent: Left: 0.13", Space After: 10 pt, Line spacing: Multiple 1.15 li, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Arial

Formatted: Justified, Indent: Left: 0.13"

Action Item 7.2 Provide maps and preserve brochures for the public in rainproof box on site.

<u>BA brochures and trail maps for the preserve</u> outlining the native plant communities, and wildlife present, and trail locations will at the preserve will be created by County staff and will be offered to visitors during the check in process. Trail maps may also be available at the specific trailheads. kept in rainproof boxes near the preserve entrance. The preserve manager or park ranger will inspect these boxes monthly and will refill the brochures as necessary.

GOAL 8: PROVIDE A PLAN FOR SECURITY AND DISASTER PREPAREDNESS

<u>Action Item 8.1</u> Discourage any unauthorized visitation to the preserve at night and identify the hours of operation.

A security light and sign designating park hours as sunrise to sunset has been will be installed at the entrances to the preserve and adjacent landowners will be given an emergency phone number if they detect human activity on the preserve after hours. If problems arise, the Collier County Sheriff's Office and/or FWC currently will be contacted to patrol the area and site on a routine basis. Once any type of camping facilities are open to the public, Aen automatic gate or temporary keypad combination lock at the entrance will allows nighttime access to the preserve to registered campers, law enforcement and staff only. Campers will be advised that the northern public use area is only available for access from dawn to dusk.

-Conservation Collier staff is developing a Preserve Ordinance which may address the issues of types of visitation, hours of operation, etc.

A single wide manufactured trailer home will be placed on a small improved area in the south public access area. A County Sheriff's Department officer or FWCC officer with jurisdictional arrest capability will reside there to keep watch over the preserve.

<u>Action Item 8.2</u> Enforce regulations prohibiting trash and landscape debris dumping in or near the preserve.

Currently, illegal dumping is not occurring on or near the preserve. Monthly property inspections will be conducted to monitor for such activity. Staff will work with the Collier County Sheriff's Office if problems start to arise.

<u>Action Item 8.3</u> Survey trees along the trail and the perimeter of the property for damage.

Staff will routinely monitor the trees along the walking and hiking trails to determine if diseased, weak, or damaged trees/limbs exist and if so remove them to reduce the risk of visitor injury. Due to the length of the proposed trails at Pepper Ranch Preserve this activity will likely require the assistance from volunteers and/or the Department of Corrections work crews, as feasible.

Formatted: Indent: Left: 0"

Formatted: Highlight

<u>Action Item 8.4</u> Visit the preserve within 48 hours after a major storm event to assess damage.

Staff will take photos of damage and fill out appropriate Collier County Risk Management Department or FEMA forms. If damage is extensive, the entire preserve or the affected portions will be closed until public safety hazards are cleared.

Action Item 8.5 Promptly clear storm debris from preserve.

If necessary, a Collier County emergency debris removal contractor will be contracted as soon as possible after the storm to schedule clean up. Removal of debris and damaged or downed trees along the trail system may be needed. Downed trees and limbs that do not appear to be a public safety hazard will be cleared at the discretion of the Preserve Manager. As much of the hurricane debris as possible will be chipped and retained onsite to be used as mulch for the trails. The preserve will be closed temporarily until the potential hazards are eliminated.

Action Item 8.6 Public Safety Response

Visitors will be given the phone number to the visitor's center or the onsite Security Officer (if after hours) incase of emergency. If a visitor is lost or injured, staff will notify the Facilities Management Department and the CCSO and EMS, will be called if needed. Staff will attempt to locate the visitor in distress by use of a 4x4 vehicle or UTV until additional help arrives. An AED (automatic electronic defibulator) is installed in the visitors center to utilize until EMS arrives. A helicopter can land close to the visitor center and the mountain bike trailhead parking area if needed. Designated landing spots will be mapped and given to EMS for future use.

GOAL 9: PROVIDE PRELIMINARY PANTHER HABITAT UNIT (PHU) CALCULATIONS AND A DRAFT MONITORING PLAN PER USFWS REQUIREMENTS FOR AN ONSITE PANTHER CONSERVATION BANK

A panther conservation bank on a portion of Pepper Ranch Preserve would create an inter-departmental partnership for Collier County. The mitigation or PHU's generated by restoring altered communities at the preserve the execution of a Conservation Easement held by a non-profit third party would be utilized for offsetting the panther impacts from Collier County transportation and other public works projects while providing the funding necessary to restore-manage the preserve.

The information provided below in Action Items 9.1 and 9.2 is meant to provide the basis necessary to allow for the commencement of preliminary discussions with all stakeholders involved in a potential future bank. If Conservation Collier and other Collier County departments choose to pursue the option of a bank further, a consultant would likely be hired to assist with planning, agency coordination and permitting of the bank.

Action Item 9.1 Provide Panther Habitat Unit calculations for the area of Pepper Ranch Preserve proposed for a Panther Conservation Bank.

Table 11 provides the calculations for the panther habitat units (PHU) for for the entire-Pepper Ranch Preserve lands that are outside of, including the areas that are within the Stewardship Sending Area 7 (SSA 7). The calculations are based on baseline the existing conditions at the preserve using current (January September 20120) USFWS habitat suitability scores.

Formatted: Not Highlight

Formatted: Not Highlight

Table 11: Panther Habitat Unit Calculations including SSA-7										
FWS Ecosyatoms	Habitat Suitability Score	Total Aeres	Panther Habitat Units							
Improved Pasture	5.2	667.34	3470.17							
Urban	0	26.26	0.00							
Dry Prairie	6.3	37.99	239.3 4							
Shrub Swamp/Brush	5.5	264.27	1453.49							
Upland Hydric Pine Forest	9.5	209.59	1991.11							
Exotic Plants	3	133.38	40.14							
Upland Hardwood Forest	9	17.42	156.78							
Hardwood Pine Forest	9.3	440.38	4095.53							
Berms	5	1.24	6.20							
Barren (Primitive Trails)	3	4.9	14.70							
Open Water	0	36.58	0.00							
Hardwood Swamp	9	241.76	2175.84							
Cypress Swamp	9	119.9	1079.10							
Marsh/Wet Prairie	4.7	309	1452.30							
		2510.01	16534.69							

To date, the USFWS has indicated to Conservation Collier staff that the SSA 7 areas will not be eligible for PHU's. For this reason, Table 12 was created to provide the total PHU's based on existing conditions excluding SSA 7.

Table-12: Panther Habita	at Unit Calcu	lations exclu	nding SSA-7						
FWS Ecosystems			Panther Habitat Units						
Improved Pasture	5.2	556.52	2893.90						
Urban	0	13.83	0.00						
Dry Prairie	6.3	36.24	228.31						
Shrub Swamp/Brush	5.5	18.68	102.74						
Upland-Hydric Pine Forest	9.5	197.59	1877.11						
Exotic Plants	3	85.42	256.26						
Upland Hardwood Forest	9	3.56	32.04						
Hardwood Pine Forest	9.3	247.21	2299.05						
Berms	5-	2.27	11.35						
Barren (Primitive Trails)	3-	2.12	6.36						
Open Water	0	27.18	0.00						
Hardwood Swamp	9	106.22	955.98						
Cypress Swamp	9	0.72	6.48						
Marsh/Wet Prairie	4.7	222.97	1047.96						
		1520.53	9717.54						

Ta
ble
11:
Pan
the
r
Ha
bita

<u>Unit Calculations excluding SSA 7 – Pre Restoration</u>

USFWS Habitat Type	FLUCFCS Code	FLUCFCS Description	Area (acres)	USFWS Assigned PHU Value	PHU's
Pine forest	411	Pine flatwoods	149.89	9.5	1,423.95
Hardwood-Pine	434	Oak, slash pine, cabbage palm	180.35	9.3	1,677.26
Cypress swamp	621	Cypress	0.72	9.2	6.64
	630	Wetland forested mix	23.81	9	214.31
Llardura ad aurama	6151	Red maple swamp	76.47	9	688.26
Hardwood swamp	6152	Pop ash swamp	2.69	9	24.21
	6162	Pond apple depression	0.71	9	6.41
Shrub swamp/brush	631	Shrub wetland	4.38	5.5	24.07
Improved pasture	211	Improved pasture	549.67	5.2	2,858.28
M 1444 B ::	641	Freshwater marsh	29.55	4.7	138.90
Marsh/Wet Prairie	643	Wet prairie	2.10	4.7	9.85
	743	Spoil	1.24	3	3.72
	3109	Upland prairie, disturbed	3.73	3	11.19
	3209	Upland shrub, disturbed	3.56	3	10.69
	4119	Pine flatwoods, distrubed	40.80	3	122.39
	4349	Oak, slash pine, cabbage palm, disturbed	0.002	3	0.01
	6189	Willow/shrub wetland, distrubed	5.44	3	16.31
	6419	Freshwater marsh, disturbed	27.21	3	81.64
	8146	Primitive trail	1.51	3	4.53
	4119E1	Pine flatwoods, disturbed, exotics 5-24%	3.39	3	10.18
	4349E1	Oak, slash pine, cabbage palm, disturbed, exotics 5- 24%	68.73	3	206.19
Barren/Disturbed Lands	6169E1	Pond apple, laurel oak, cabbage palm, disturbed, exotics 5-24%	0.83	3	2.49
	6319E1	Shrub wetland, disturbed, exotics 5-24%	4.72	3	14.17
	6419E1	Freshwater marsh, disturbed, exotics 5-24%	70.00	3	210.00
	4119E2	Pine flatwoods, disturbed, exotics 25-49%	1.24	3	3.71
	4349E2	Oak, slash pine, cabbage palm, disturbed, exotics 25-49%	8.86	3	26.57
	6319E2	Shrub wetland, disturbed, exotics 25-49%	0.09	3	0.26
	6419E2	Freshwater marsh, disturbed, exotics 25-49%	26.19	3	78.57
	6439E2	Wet prairie, disturbed, exotics 25-49%	7.50	3	22.50
	6419E3	Freshwater marsh, disturbed, exotics 50-74%	3.46	3	10.37
	743E4	Spoil, exotics 75-100%	8.16	3	24.47
	4119E4	Pine flatwoods, disturbed, exotics 75-100%	3.69	3	11.06

USFWS Habitat Type	FLUCFCS Code	FLUCFCS Description	Area (acres)	USFWS Assigned PHU Value	PHU's
	422	Brazilian pepper, non-hydric	0.40	3	1.20
	6192	Brazilian pepper, hydric	0.84	3	2.53
Exotic/Nuisance Plants	437	Australian pine	1.16	3	3.47
	428E1	Cabbage palm, exotics 5-24%	1.14	3	3.41
	428E3	Cabbage palm, exotics 50-74%	3.29	3	9.86
	180	Campground	11.86	0	-
Urban	700	Cattle Dipping Vat Remediation Area	1.02	0	-
	8145	Shell road, graded and drained	14.31	0	-
	512	Ditches	24.92	0	-
Water	512E4	Ditches, exotics 75-100%	0.66	0	-
	742	Borrow pond	0.78	0	-
Dry prairie with 14.5% exotic plant coverage	310E1	Upland prairie, exotics 5-24%	35.11	6.3/3	204.42
Dry prairie with 37% exotic plant coverage	310E2	Upland prairie, exotics 25-49%	4.01	6.3/3	20.36
Hardwood Forest with 14.5% exotic plant coverage	427E1	Oaks, exotics 5-24%	1.57	9/3	12.73
Hardwood Swamp with 14.5% exotic plant coverage	630E1	Wetland forested mix, exotics 5-24%	1.80	9/3	14.61
Shrub swamp/brush with 14.5% exotic plant coverage	631E1	Shrub wetland, exotics 5-24%	1.35	5.5/3	6.91
Marsh/Wet Prairie with 14.5% exotic plant coverage	641E1	Freshwater marsh, exotics 5-24%	83.40	4.7/3	371.43
Marsh/Wet Prairie with 37% exotic plant coverage	641E2	Freshwater marsh, exotics 25-49%	17.26	4.7/3	70.27
Marsh/Wet Prairie with 62% exotic plant coverage	641E3	Freshwater marsh, exotics 50-74%	1.27	4.7/3	4.62
TOTAL			1,516.84		8,669.0

To determine the amount of PHU's that could be available for mitigation, the <u>above</u> following calculations were performed based on site conditions <u>post-pre</u> restoration. Site restoration includes converting improved/unimproved pastures, ditches, berms, and exotic communities to upland or hydric pine forest. Restoration also includes conversion of a borrow pond to marsh community. Tables 13 and 14 provide the post restoration PHU calculations including and excluding SSA 7, respectivelyUSFWS has informed County staff that credit will only be given for restoration outside of the scope of this management plan. Control of invasive, exotic vegetation and prescribed fire will not

Pepper Ranch Preserve Land Management Plan

result in additional PHU credits. The Preserve, excluding SSA 7, could provide approximately 8,669.0 PHUs. These preliminary PHU calculations are based on current (September 2012) USFWS habitat suitability scores and are subject to review and final approval by USFWS.

Formatted: Indent: Left: 0"

	Habitut Suitability Score	Total Acres	Panther-Habitat Units
Improved Pasture to Upland-Hydric Pine Forest	7.35	*640.74	4709.44
Urban	0	26.26	0.00
Dry Prairie	6.3	39.45	248.54
Shrub Swamp/Brush	5.5	266.52	1465.86
Upland-Hydric Pine Forest	9.5	219.40	2084.49
Non native (exotics) to Upland Hydric Pine Forest	6.25	14.86	92.88
Upland Hardwood Forest	9	20.92	188.28
Hardwood-Pine Forest	9.3	464.19	4316.97
Barren (Primitive Trails)	3	4.90	14.70
Hardwood Swamp	9	242.46	2182.14
Cypress Swamp	9	126.07	1134.63
Marsh/Wet Prairie	4.7	367.08	1725.28
Berm Restoration to Upland-Hydric Pine Forest	7.25	2.43	17.62
Ditch to Upland Hydric Pine Forest	4.75	35.15	166.96
Borrow Pond to Marsh	4.7	1.43	6.72
		1831.12	18354.51

*Approximately 38.13 acres of various habitat will be converted to recreational uses (i.e. trailhead parking areas, RV camping, campgrounds, etc.). This acreage has been removed from the PHU calculations.

Formatted: Normal, Left, Space Before: 0 pt, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Normal, Left, Space Before: 0 pt, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted Table

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Indent: Left: 0", Right: 0"

Formatted: Left, Indent: Left: 0"

	Habitat Suitability Score	Total Acres	Panther Habitat Units
Improved Pasture to Upland-Hydric Pine Forest	7.35	<u>*529.92</u>	3894.91
Urban	0	13.83	0.00
Dry Prairie	6.3	37.39	235.56
Shrub Swamp/Brush	5.5	19.61	107.86
Upland-Hydric Pine Forest	9.5	199.52	1895.44
Non native (exotics) to Upland Hydric Pine Forest	6.25	9.96	62.25
Upland Hardwood Forest	9	6.00	54.00
Hardwood-Pine Forest	9.3	259.23	2410.84
Barren (Primitive Trails)	3	2.12	6.36
Hardwood Swamp	9	106.61	959.49
Cypress Swamp	9	0.72	6.48
Marsh/Wet Prairie	4.7	268.04	1259.79
Berm Restoration to Upland-Hydric Pine Forest	7.25	2.27	16.46
Ditch to Upland Hydric Pine Forest	4.75	26.40	125.40
Borrow Pond to Marsh	4.7	0.78	3.67
		1482.40	11038.51

^{*}Approximately 38.13 acres of various habitat will be converted to recreational uses (i.e. trailhead parking areas, RV camping, campgrounds, etc.). This acreage has been removed from the PHU calculations.

Subtracting the total PHUs pre restoration from the total PHUs post restoration provides the "lift" or PHUs available for mitigation. Based on existing and post restoration conditions at the Pepper Ranch Preserve, the entire Preserve (including SSA 7) could provide approximately 1,819.82 PHUs. The Preserve, excluding SSA 7, could provide approximately 1,320.97 PHUs. The preliminary PHU calculations are based on current (January 2010) USFWS habitat suitability scores and are subject to review and final approval by USFWS.

Action Item 9.2 Provide a draft Monitoring Plan per USFWS requirements for the area of Pepper Ranch Preserve proposed for a Panther Conservation Bank.

With the establishment of a panther conservation bank the USFWS requires a monitoring plan for the lands within the designated bank to ensure the bank continues to meet its success criteria in perpetuity. Below is a draft monitoring plan for the creation of a possible panther conservation bank at Pepper Ranch Preserve.

Formatted: Normal, Left, Space Before: 0 pt, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Normal, Left, Space Before: 0 pt, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left. Don't adjust space between Latin and Asian

text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian

text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian

text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian

text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Left, Indent: Left: 0", Right: 0"

Formatted: Left

Formatted: Left, Indent: Left: 0"

Monitoring

Baseline monitoring will be completed by a consultant within 60 days of approval of the Bank by the Service and a baseline monitoring report will be forwarded to Service staff in Vero Beach within 45 days of the monitoring event. Time-zero monitoring will be completed within 60 days of the completion of initial prescribed fires. As with the baseline monitoring report, the time-zero monitoring report will be forwarded to Service staff in Vero Beach within 45 days of the monitoring event. Annual monitoring will begin 12 months following the time-zero monitoring event and continue for a total of five years. Annual monitoring reports will be forwarded to Service staff in Vero Beach prior to January 31 each year. If, at the end of five years of monitoring, the Bank has reached success criteria, monitoring will be conducted once every five years to ensure that success criteria are met in perpetuity. If success criteria are not met, annual monitoring will continue until they are achieved. A summary of the reporting schedule can be found in Baseline monitoring will be completed within 60 days of approval of the Bank by USFWS and a baseline monitoring report will be forwarded to USFWS staff in Vero Beach within 45 days of the monitoring event. Time zero monitoring will be completed within 60 days of the completion of exotic and nuisance vegetation removal. As with the baseline monitoring report, the time-zero monitoring report will be forwarded to USFWS staff in Vero Beach within 45 days of the monitoring event. Annual monitoring will begin 12 months following the time zero monitoring event and continue for a total of five years. Annual monitoring reports will be forwarded to USFWS staff in Vero Beach prior to January 31 each year. If, at the end of five years of monitoring, the Bank has reached success criteria, monitoring will be conducted once every five years to insure that success criteria are met in perpetuity. If success criteria are not met then annual monitoring will continue until met. A summary of the reporting schedule can be found in Table 15.

Table 3332: Monitoring and Reporting Schedule for Panther Conservation Bank

Report	Monitoring Implemented	Delivery
Baseline Monitoring	Within 60 Days of Approval	45 days
Time-Zero Monitoring	Within 60 Days of Initial Restoration	45 days
Annual Monitoring	Year 1 12 Months After Time-Zero	45 days
_	Monitoring	
Annual Monitoring	Year 2 1 Year from Previous Report	January 31
Annual Monitoring	Year 3 1 Year from Previous Report	January 31
Annual Monitoring	Year 4 1 Year from Previous Report	January 31
Annual Monitoring	Year 5 1 Year from Previous Report	January 31
Five-Year Monitoring	5 Years from Previous Report	January 31
(Year 10)	•	
Every 5 years thereafter	5 Years from Previous Report	January 31

In addition to the information outlined below, the monitoring report will include a general overview of the land management activities (i.e. prescribed burns, exotic vegetation maintenance, pasture restoration activities, etc.) conducted since the

Formatted: Indent: Left: 0.13"

Formatted: Font: Not Bold

Formatted: Font: Not Bold

previous monitoring report and planned maintenance and management activities during the next period.

Vegetation Monitoring:

Permanent monitoring transects will be established during the baseline monitoring event and located throughout the site to include a thorough representation of the various habitats onsite. Three vegetative strata will be sampled along each transect and will be representative of habitat types throughout the site. These strata are: overstory [plants greater than four inches diameter breast height (DBH)], understory (plants greater than four inches DBH and greater than three feet in height), and ground cover (all non-woody plants and woody plants less than three feet in height). The overstory and understory vegetation will be sampled in 10 m2 plots and the ground cover vegetation will be sampled in 1m2 plots along each monitoring transect. Panoramic photographs will be taken at the beginning of each transect to provide physical documentation of the condition and appearance of the property as well as any changes taking place. The panoramic photographs will be included in each monitoring report. For the overstory and understory strata, the relative canopy closure for each species will be recorded. Average shrub height will be recorded for all species identified in the understory stratum. Percent coverage and average height for all saw palmetto will be recorded for plots located within habitats with saw palmetto. The percent cover of groundcover species and bare ground will be estimated for the herbaceous study plots along each transect. Exotic and nuisance vegetation coverage within the plots will be recorded. Survival rate evaluations will occur throughout the site to include a thorough representation of the various habitats onsite. There will be a maximum of 17 transects with a total of 51 sample plots.

Permanent monitoring transects will be established during the time zero monitoring event and located throughout the site to include a thorough representation of the various natural communities onsite. Three vegetative strata will be sampled along each transect and will be representative of natural community types throughout the site. These strata are: canopy [plants >4 inches diameter breast height (DBH)], midstory (plants <4 inches DBH and greater than 3 feet in height) and ground cover (all non-woody plants and woody plants less than 3 feet in height). The canopy and midcanopy vegetation will be sampled in 10 square meter plots and the ground cover vegetation sampled in 1 square meter plots along each monitoring transect. Panoramic photographs will be taken at the beginning of each transect to provide physical documentation of the condition and appearance of the property as well as any changes taking place. The panoramic photographs will be included in each monitoring report. For the canopy and midcanopy stratums, the relative canopy closure for each species will be recorded. Average shrub height will be recorded for all species identified in the midcanopy stratum. Percent coverage and average height for all saw palmetto will be recorded for plots located within communities with saw palmetto. The percent cover of ground cover species and bare ground will be estimated for the herbaceous study plots along the transects. Exotic and nuisance vegetation coverage within the plots will be recorded. Survival rate evaluations will occur throughout the site to include a thorough representation of the various natural communities onsite.

Exotic and Nuisance Species Monitoring:

In addition to the permanent monitoring transects, existing disturbed areas, such as fence lines, fire breaks, and primitive roads / trails, will be surveyed annually, using the FWC protocol, by vehicle and meandering pedestrian transects to assess the site for the presence and percent coverage of exotic vegetation species. Following the annual exotic vegetation surveys, an exotic vegetation map will be prepared illustrating the locations of exotic and nuisance vegetation in need of corrective action. The map will be provided to the County contractor annually to ensure timely and effective treatment. In addition to the permanent monitoring transects, existing disturbed areas, such as fence lines, fire breaks and primitive roads/trails, will be surveyed annually, using the FWC protocol (FWC 2009), by vehicle and meandering pedestrian transects to assess the site for the presence and percent coverage of exotic vegetation species. Following the annual exotic vegetation surveys, an exotic vegetation map will be prepared illustrating the locations of exotic and nuisance vegetation in need of corrective action. The map will be provided to the County and USFWS annually to ensure timely and effective treatment.

Wildlife Utilization:

Spotlight transect surveys will be utilized to census white-tailed deer due to the large acreage of open habitat within the Preserve, density of forested habitat, and the available roads and trails. For each transect the spotlighting visibility will be estimated once per season, before conducting the spotlight census. The spotlighting visibility will be calculated as the acreage of habitat perpendicular to each transect which can be surveyed for white-tailed deer. Visibility will be dependent on the density and height of vegetation and also the terrain. Two hundred yards will be the maximum distance from which visibility will be quantified and white-tailed deer will be censused. Visibility stations will be placed every 0.10 miles along and at the beginning and end of each transect. At each visibility station a one-million candle power spotlight will be used to illuminate the habitat perpendicular to both sides of each transect. A Bushnell Laser Range Finder Sport 450 will be used to determine the distance, in yards, to the nearest obstruction which would deter viewing a deer on either side of each transect. The laser range finder will have an accuracy of +/- one yard. The spotlighting visibility per transect and cumulative spotlighting visibility will be calculated as the acreage of visibility.

Spotlight transect surveys will begin one-half hour after sunset. The deer spotlight census will follow the methodology described by Mitchell (1986). Six transects will be surveyed in order during each census and each transect will be surveyed without interruption until completed. All census data for each transect will be recorded on a separate data sheet. The data recorded will include: transect number, official sunset, date, time survey began, time survey ended, temperature, wind direction, average wind speed, percent cloud cover, name of personnel, number of bucks, number of does, number of fawns, and number of unknown deer.

A minimum of four personnel will be utilized for each of the spotlighting censuses: one driver, one data recorder, and two spotlight observers. A four-wheel drive pickup truck will be utilized for each census and the spotlight observers will be stationed in the bed of the pickup. For each transect the vehicle will be driven at 5-10 mph and each spotlight observer will scan the habitat on their side of the vehicle with a one-million candle power

spotlight. If a deer is observed the vehicle will briefly stopp and the spotlight observer will use binoculars to identify the age and sex of each deer observed.

For each group of deer the spotlight observers will classify each deer as either buck, doe, fawn, or unidentified. A group will consist of one single deer by itself or more than one deer grouped together; and the grouping of deer will be subjective - meaning the spotlight observer will determine how deer in an area are grouped. Sex and age will be recorded for each deer only if all the deer in that group can be sexed and aged. If one deer in the group cannot be identified, then all the deer in the group will be classified as unidentified in order to reduce bias when estimating the total number of bucks, does, and fawns on the Preserve.

The annual wildlife monitoring reports will include the following information:

- Results of the annual spotlight survey.
- A brief description of work performed since the previous report (if applicable) along with a discussion of any modifications to the survey methodology.
- A list of all wildlife species observed during the survey.
- Direct evidence (i.e., tracks, scat, visual sightings, and rub trees) of panther preyspecies observed during each sampling period.
- Hunt harvest data (if applicable).

Regular and periodic observations of wildlife will be made during all monitoring events and other site visits by qualified ecologists. This will consist of recording evidence and signs of wildlife (i.e., direct sightings, vocalizations, burrows, nests, tracks, droppings, etc.). The number of white tailed deer, feral hog, and panther observations at the site will be recorded during each monitoring event and included in the annual reports. Regular and periodic observations of wildlife will be made during all monitoring events and other site visits by qualified ecologists. This will consist of recording evidence and signs of wildlife (i.e., direct sightings, vocalizations, burrows, nests, tracks, droppings, etc.). The number of white tailed deer, feral hog and panther observations at the site will be recorded during each monitoring event and included in the annual reports. In addition, permanent wildlife transects will be established in representative natural communities onsite from which surveys for panther and panther prey tracks will be conducted. Efforts will be made to establish the wildlife transects in areas that are conducive to track detection and identification with as little disturbance by vehicular traffic as possible. The wildlife transects will be monitored annually for a 5 day period, and results submitted in conjunction with the vegetation monitoring in the monitoring report. Updated panther radio telemetry data (when available) within a 2 mile radius of the site will be included in the monitoring report to document radio collared panther activity in the Bank vicinity over the course of the monitoring period.

In addition to the information outlined above, the monitoring report will also include a general overview of the land management activities (i.e. prescribed burns, exotic vegetation maintenance, pasture restoration activities, etc.) conducted since the previous monitoring report and planned maintenance and management activities during the next period.

Formatted: Indent: First line: 0.5"

Formatted: Indent: Left: 0.5"

Formatted: Indent: First line: 0.5"

Formatted: Indent: Left: 0.5"

Formatted: Indent: First line: 0.5"

4.5 Establish an Operational Plan for the Pepper Ranch Preserve

This section provides management recommendations for operation of the Pepper Ranch Preserve. It discusses maintenance and budgeting needs, the possibilities for contracting the restoration activities, coordination and other management issues.

4.5.1 Maintenance

Initially, the primary maintenance activities for the preserve will includes invasive exotic species control and trail maintenance and site security. Particularly important are the security measures to prevent trespassingkeep intruders out and to maintain the signage and fencing (where installed) in good condition. Signs that effectively convey the desired message provide an opportunity for increasing environmental education and awareness. Once the preserve is open to the public other sSignificant maintenance activities will be necessary for the upkeep of all public facilities including but not limited to the trailheads/parking areas, visitor centerlodge, campgrounds, boardwalks, restrooms and interpretive signage.

4.5.2 Estimated Annual Costs and Funding Sources

Preliminary budget estimates for Pepper Ranch Preserve include cost breakdowns associated with resource restoration and management. The funding source identified for the restoration and management activities is the Conservation Collier Program Management Trust Fund. Grants will be sought to supplement existing management funds particularly for the areas within SSA 7. Staff will also utilizes the Collier County Sheriff's Department weekend workers programs and the Civil Citation program for certain labor projects and may also separately involve the County Scout programs and volunteers for trail maintenanceereation and enhancement.

The budget in Table 146-13 represents the actual and unmet budgetary needs for managing the lands and resources of the preserve overfor the next ten years. The table shows the actual costs of land management activities, construction costs since acquisition and the estimated costs over the next 3 years. the activities planned and the initial and annual cost estimate of each activity. The assumption was made that public facilities at the south public access area would be constructed first. This budget was developed using data from Conservation Collier and other cooperating entities, and is based on actual costs for land management activities, equipment purchase and maintenance, and for development of fixed capital facilities. The budget considers available funding and is consistent with the direction necessary to achieve the goals and objectives for Pepper Ranch Preserve.

In August of 2017, a thorough building assessment was conducted by engineers and building inspectors on all the structures on the property. It was determined that several structural issues need to be addressed to maintain the historical visitor center and pole barns. Maintenance on these structures has been budgeted for in the fical year 2017-18. However, after much deliberation, staff has decided that the cottage/caretaker's home will have to be torn down. It has had major structural issues since we purchased the

property, and the cost to continue to mitigate these ongoing problems will be excessive. A concrete pad will be installed at the north end of the campground with a full hook-up to allow a campground host to reside there during the months that the preserve is open. Decisions will have to be made in the future on whether or not to also install a manufactured home on the property for an enforcement officer to live in.

There are presently no utilities on site; a septic system and well for potable water will need to be installed. Determining the extent and capacity of public use facilities to be developed will help determine the type and amount of utilities required. Due to these current unknowns it is difficult to estimate the cost of both the utilities and the public use facilities. The cost estimates provided in Table 16 are based on best available knowledge, are subject to change and many could not be given at this time.

			7P-1-1-	16. 10.41	4							
			Table	16: Esti	muted A	nnual La	nd Mana ;	gement B	udget	<u> </u>		
							2014-15				2019-20	Total
Facilities Development												
Permit, planning, design ¹			\$90,000									\$90,000
Permit application fees ^{1,2}	2		\$ 2,250									\$2,250
Entrance gate ³	1			\$ 25,000								\$25,000
Interpretive signs ⁴	2	\$500		\$ 500			\$ 500					\$1,000
Plant signs ⁵	50	\$10		\$ 250			\$ 250					\$500
Entrance signage ⁶	2	\$2,000		\$ 4,000								\$4,000
Directional signage ⁷	2	\$200		\$ 400								\$400
Benches ⁸	6	\$650		\$ 1,300	\$ 2,600							\$3,900
Check in/gate booth ^{1,9}				\$ 10,000								\$10,000
Electrical Power Supply ^{1,10}				\$-30,000								\$30,000
Site, fill and mise. 1,11				\$ 50.000								\$50,000
Walkways ^{1,12}	500 LF	\$20/LF		\$ 10,000								\$10,000
Boardwalk ^{1,13}	250 LF	\$350/LF					\$87,500					\$87,500
Observation Platform ^{1,14}							\$ 8,000					\$8,000
Driveway into lodge^{1,15}	8,000 ft ²	\$5/ft ²					\$ 40,000					\$40,000
Parking area 16	1			\$ 30,000								\$30,000
Septic system for trailer				\$ 40,000								\$40,000
Septic system for restrooms ¹				\$ 75,000								\$75,000
Well and treatment system for trailer ¹				\$-30,000								\$30,000
Well and treatment system for restrooms ¹				\$ 75,000								\$75,000
Restroom/shower building ¹				\$144,000								\$144,000
Caretaker house ^{1,17}				\$ 3,000								\$3,000

Water W. Water	4 - 3 4		136										
·1 able 16: Esuma	tea An n	iuai Lanc	l Manage	ment-Bud	g et (conti	nuea)					l		
	QTY		2010-11	2011-12	2012-13			2015-16					
Temporary portable toilets (1 ADA; 1 regular; weekly maintenance) ¹⁸	6 months	\$290/ month	\$ 1,740										\$ 1,740
Ramps to lodge and trailer ¹	2	\$2,000		\$ 4,000									\$ 4,000
Installation of trailer for onsite law enforcement ¹⁹				\$ 50,000									\$ 50,000
Restoration/Monitoring													
Firebreak and trail clearing/mowing ²⁰	30,000 1.f.	\$3.33/ LF	\$100,000	\$ 25,000	\$15,000	\$10,000	\$10,000	\$-5,000	\$-5,000	\$ 5,000	\$-5,000	\$-5,000	\$ 185,000
Establish photo points	10		\$ 150	\$ 30	\$ 30	\$ 30	\$ 30	\$ 30	\$ 30	\$ 30	\$ 30	\$ 30	\$ 420
Remove invasive exotic and nuisance species			\$240,000	\$500,000	\$97,500	\$97,500	\$72,500	\$ 72,500	\$72,500	\$72,500	\$72,500	\$72,500	\$1,370,000
Mitigation													
Panther/wetland eonservation bank planning and permitting		\$91,000	\$50,000	\$ 41,000									\$-91,000
Vegetation Monitoring					\$50,000	\$17,000	\$17,000	\$17,000	\$17,000	\$17,000	\$17,000	\$17,000	\$169,000
Regular Maintenance						·		·			·		
Preserve assist (seasonal)				\$ 15,000	\$15,000	\$15,000	\$15,000	\$15,000	\$15,000	\$15,000	\$15,000	\$15,000	\$135,000
Lawn mowing			\$ 4,000	\$ 4,000	\$ 4,000	\$ 4,000	\$ 4,000	\$ 4,000	\$ 4,000	\$ 4,000	\$ 4,000	\$ 4,000	\$ 40,000
Electricity			\$ 1,200	\$ 1,200	\$ 1,200	\$ 1,200	\$ 1,200	\$ 1,200	\$ 1,200	\$ 1,200	\$ 1,200	\$ 1,200	\$ 12,000
Cleaning service			\$ 400	\$ 400	\$ 400	\$ 400	\$ 400	\$ 400	\$ 400	\$ 400	\$ 400	\$ 400	\$ 4,000
Misc. Facility Repairs			\$10,000	\$ 5,000	\$-5,000	\$ 5,000	\$ 5,000	\$-5,000	\$ 5,000	\$ 5,000	\$ 5,000	\$ 5,000	\$ 55,000
Brochures			\$ 300	\$ 300	\$ 300	\$ 300	\$ 300	\$ 300	\$ 300	\$ 300	\$ 300	\$ 300	\$ 3,000
GRAND TOTAL:			\$500,040	\$1,174,380	\$191,030	\$150,430	\$261,680	\$120,430	\$120,430	\$120,430	\$120,430	\$120,430	\$2,879,710

Formatted: Left

Assumptions for Cost Estimates:

- 1. CH2MHill Project costs
- 2. Permit application fees: SFWMD Standard General Permit \$1,750; Collier DOH \$500
- 3. Entrance gates: Two electronic gates, one at each of the entrances to the preserve, that would provide nighttime access to campers, law enforcement and preserve staff.
- 4. **Interpretive signs:** 2 interpretive signs near the lodge, 4'x6' at \$500 each.
- 5. Plant signs: small signs identifying native plants, 50 @ \$10 each.
- 6. Entrance signage: 2 arched "ranch style" entrance signs over both entrances estimated @ \$2,000 each.
- 7. **Directional signage:** 2 road signs indicating entrance to the preserve—1 on SR29 and 1 on Lake Trafford Road (\$200 each).
- 8. Benches: each bench at \$650; benches could be constructed as part of Eagle Scout projects to eliminate this cost.
- 9. Check in/gate booth: located at north entrance with power, AC and phone.
- 10. Electrical power supply: supply to lighting near campground and restroom building.
- 11. Site, fill, mise.: Cost to fill bathroom building area and driveway widening.
- 12. Walkways: 5' wide; pervious
- 13. Boardwalk: eeo-friendly, 6' wide or 4' wide with handicapped turn arounds.
- 14. Observation Platform: Roofed area at end of boardwalk over looking lake 20° X 20°

- 15. Driveway to lodge: 500 L.F. X 16' wide; culverts may need to be installed for stormwater purposes.
- 16. Parking areas: 1 parking area for up to 30 pervious spaces (\$30,000)
- 17. Caretaker house: gut and convert to storage
- 18. Temporary portable toilets: \$290/month for 6 months, based on United Site Services quote from IMP; this is a possible option until permanent facilities are constructed.
- 19. Installation of trailer for onsite law enforcement: assumes a brand new single wide trailer.
- 20. Firebreak and trail clearing/mowing: based upon estimates from Collier County contractor in similar habitats.

Formatted: No bullets or numbering

Table 16: Estimated Annual Land Management Budget												
Item	2009-10	2010-11	2011-12	2012-13	2013-14	2014-15	2015-16	2016-17	2017-18	2018-19	2019-20	Total
BROCHURES				\$102	\$600	\$85		\$85	\$200	\$200	\$200	\$ 1,472
CONSTRUCTION/IMPROVEMENTS	\$6,578	\$85,699		\$3,445	\$18,824	\$115,752	\$16,592	\$1,365	\$74,000			\$322,255
CONSULTING/SURVEYS	\$60,653	\$11,287		\$7,350		\$2,400	\$17,800	\$15,840				\$115,330
COUNTY DEPT. SERVICES	\$12,229	\$10,373	\$1,979	\$1,400		\$797		\$448	\$500	\$1,000	\$1,000	\$29,727
DEBRIS SERVICES	\$4,770				\$9,070							\$13,840
EMERGENCY SERVICES					\$1,267	\$1,557	\$600					\$3,424
FIREBREAK MOWING TRAILS		× ×		SZ				\$8,200	\$8,000	\$5,000	\$5,000	\$26,200
EXOTIC MAINTENANCE	\$41,000	\$255,627	\$99,904	\$67,800	\$245,454	\$182,128	\$128,683	\$137,034	\$104,800	\$105,000	\$105,000	\$1,472,430
FENCING				\$3,550.00								\$3,550.00
FIELD SUPPLIES & EQUIPMENT	\$1,613	\$50.59	\$882	\$361	\$5,334	\$967	\$333					\$9,540
LABOR SERVICES	\$756	\$2,431.50	\$111,742	\$3,600	\$3,455	\$1,500	\$1,500.00	\$1,000	\$1,000	1,000	\$1,000	\$128,985
LANDSCAPING SERVICES	\$9,678	6. 50 0. 50		\$8,180	\$1,273	\$10,100	\$7,500.00	\$5,485	\$500	5,000	\$500	\$48,216
LEASE/LEGAL					\$616		\$512.46		\$500	\$500	\$500	\$2,628
LICENSING & PERMITTING	\$91,040		\$30,688	\$1,844	, , , , , , , , , , , , , , , , , , ,	\$1,467	\$50.00		100	100	100	\$125,388
MONITORING		× ×		SZ VI	\$900							\$900
MOWING (ALL)	\$1,678	\$34,990	\$8,890	\$5,640.00			100					\$51,198
JANITORIAL				\$428	\$2,892		\$44		\$ 1,000	\$1,000	\$1,000	\$6,364
OFFICE SUPPLIES	\$1,599		\$372	\$460	\$841	\$58		\$238	\$200	\$200	\$200	\$4,168
PEST ABATEMENT					\$847			\$625	\$500	\$500	\$500	\$2,972
PORT-A-POTTY	\$965	\$1,390	\$2,750	\$2,415	\$500	\$645.03	\$460	\$460	\$500	\$500	\$500	\$11,085
SIGNS	\$417		\$166	\$464	\$601			\$362	\$200	\$200	\$200	\$2,610
TAXES	. (20)	\$163	\$166	\$192	\$160	\$164	\$170	\$176	\$180	\$184	\$190	\$1,745
UTILITIES (ALL)	\$993	\$1,400	\$940	\$2,010.00	\$3,968	\$5,236	\$5,530	\$6,163	\$6,400	\$6,400	\$6,400	\$45,441
VOLUNTEER RELATED						\$214.57		\$27	\$100	\$100	\$100	\$ 541.77
GRAND TOTAL:	\$ 233,969	\$403,411	\$258,478	\$109,240	\$296,601	\$323,071	\$179,774	\$ 177,508	\$198,680	\$126,884	\$122,390	\$2,430,008

^{*}Actual numbers are included from FY 2009-10 through 2016-17. Numbers are estimated FY 2017-18 through 2019-2020.

^{*}Labor Services: Fees associated with Check Station Attendant for Public Hunts
*Utilities: include electricity, water, garbage service and phone and internet

4.5.3 Potential for Contracting Restoration and Management Activities by Private Vendors

A significant number of management operations and restoration activities on the Pepper Ranch Preserve can be considered for outsourcing. Restoration and management activities that can be considered for outsourcing to private entities are listed in Table $\frac{1014}{1}$.

Table 1912: Potential Contracting for Restoration and Management Activities									
Activity	Approved	Conditional	Rejected						
Prescribed fire and/ or mechanical treatment application	X								
Minor fireline installation	X								
Fireline, fence and trail maintenance	X								
Fence installation	X								
Plant and wildlife inventory and monitoring		X							
Listed species mapping and needs assessment		X							
Restore/enhance encroachment and ruderal areas		X							
Reduce exotic species	X								
Literature development and printing		X							
Interpretive signs development and installation		X							
Trail installation	X								
Parking Area construction	X								
Law enforcement and patrol	X								

5.0 LITERATURE CITED

- Abrahamson, W. G., and D. C Hartnett. 1990. Pine flatwoods and dry prairies. Pages 103-149 in R. L. Myers and J. J. Ewel editors. Ecosystems of Florida. University of Central Florida Press; Orlando, Florida.
- Brown, P.M. 2002. Wild Orchids of Florida. Gainesville: The University Press of Florida. 409 p.
- Bush, C.S., and J.F. Morton. 1969. Native Trees and Plants for Florida Landscaping. Pages 8-9. Florida Department of Agriculture and Consumer Services.
- Campbell K. M. 1990. Soil survey of Collier County area Florida. USDA, Natural Resources Conservation Service; Washington, D.C.
- Campbell, T. 2001. The brown anole. Institute for Biological Invaders: Invader of the Month. University of Tennessee, Knoxville, TN. Available from http://invasions.bio.utk.edu/invaders/sagrei.html (accessed November 2007).
- Campbell, T. S. 1996. Northern range expansion of the brown anole, *Anolis sagrei*, in Florida and Georgia. Herp. Review 27:155-157.
- Campbell, T. S. 2000. Analyses of the effects of an exotic lizard (*Anolis sagrei*) on a native lizard (*Anolis carolinensis*) in Florida, using islands as experimental units. Ph.D. Dissertation, University of Tennessee, Knoxville, TN.
- Coates, S. F., M. B. Main, J. J. Mullahey, J. M. Schaefer, G. W. Tanner, M. E. Sunquist, and M. D. Fanning. 1998. The coyote (*Canis latrans*): Florida's newest predator. University of Florida Cooperative Extension Service Document WEC124. 5pp. University of Florida, UF/IFAS Extension Digital Information Source (EDIS) Database. Available from http://edis.ifas.ufl.edu/pdffiles/UW/UW12700.pdf (accessed November 2007).
- Cobb, B., E. Farnsworth, C. Lowe. 2005. Ferns of Northeastern and Central North America. New York: Houghton Mifflin Company. 417 p.
- Endangered Species Act. US Code Title 16 Chapter 35 § 1532 (19) (1973). Available from http://www.fws.gov/Endangered/pdfs/esaall.pdf
- eFloras. Flora of North America. Available from http://efloras.org/florataxon.aspx?flora_id=1&taxon_id=233500308 (accessed December 2009)

- Fairbank, P. and S. Hohner. 1995. Mapping recharge (infiltration and leakage) throughout the South Florida Water Management District. Technical publication 95-20 (DRE # 327). SFWMD, West Palm Beach, Florida.
- Federal Endangered Species Act of 1973, as amended, Pub. L. No. 93-205 (87 Stat. 884).
- Florida Department of State (FDOS). 2006. Aerial Photography of Florida, a State University System of Florida PALMM Project. Available from http://web.uflib.ufl.edu/digital/collections/flap/Counties/index.html (accessed on December 2009).
- Florida Department of Transportation 1999. Florida Land Use and Cover Forms Classification System (FLUCFCS).
- Florida Exotic Pest Plant Council (FLEPPC). 2009. List of Florida's invasive plant species. Florida Exotic Pest Plant Council. Available from http://www.fleppc.org/list/09list.htm (accessed December 2009).
- Florida Fish and Wildlife Conservation Commission (FWC). 1999-2010. Species information coyote. Available from http://www.myfwc.com/WILDLIFEHABITATS/SpeciesInfo_Coyote.htm (accessed on December 2009).
- Florida Fish and Wildlife Conservation Commission (FWC). 2003. Florida's breeding bird atlas: A collaborative study of Florida's birdlife. http://www.myfwc.com/bba/(accessed December 2009).
- Florida Fish and Wildlife Conservation Commission (FWC). 2007. The Coyote in Florida: Compiled by Walter McCown and Brian Scheick. Fish and Wildlife Research Institute. Available from http://www.myfwc.com/docs/WildlifeHabitats/CoyoteWhitePaperFinal.pdf (accessed July 2010)
- Florida Fish and Wildlife Conservation Commission (FWC). 2009. Standard Exotic Plant Survey Protocol. Tallahassee, Florida.
- Florida Fish and Wildlife Conservation Commission (FWC). 2009. Florida's Endangered Species, Threatened Species, and Species of Special Concern. Available from http://www.myfwc.com/docs/WildlifeHabitats/Threatened_Endangered_Species.pdf (accessed on December 2009).
- Florida Natural Areas Inventory (FNAI) and Florida Department of Natural Resources (FDNR) 1990. Guide to the Natural Communities of Florida. Florida Natural Areas Inventory and Florida Department of Natural Resources.

- Florida Natural Areas Inventory (FNAI). Natural Communities. Tallahassee (FL). Available from: http://fnai.org/pdf/MAxCounty_201703.pdf http://www.fnai.org/descriptions.cfm (accessed AugustJanuary 20170).
- Gann, G. D., K. A. Bradley, and S. W. Woodmansee. 2002. Rare Plants of South Florida: Their History, Conservation, and Restoration. The Institute for Regional Conservation. Miami, Florida.
- Giuliano, W. M., and G. W. Tanner. 2005. Control and management of wild hogs in Florida. Department of Wildlife Ecology and Conservation Publication WEC 192. 7pp. Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences (IFAS), University of Florida. Available from http://edis.ifas.ufl.edu/UW221 (accessed December 2007).
- Giuliano, W. M., and G. W. Tanner. 2005. Ecology of wild hogs in Florida. Department of Wildlife Ecology and Conservation Publication WEC 191. 7pp. Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences (IFAS), University of Florida. Available from http://edis.ifas.ufl.edu/UW220 (accessed December 2007).
- Humphrey, S.R. 1992. Florida black bear. Pp. 265-275. In S.R. Humphrey (ed.), Rare and Endangered Biota of Florida, Mammals. University of Florida, Gainesville, FL.
- Humphrey, S.R. and P.G.R. Jodice. 1992. Big Cypress fox squirrel. Pp. 224-233. In S.R. Humphrey (ed.), Rare and Endangered Biota of Florida, Mammals. University of Florida, Gainesville, FL.
- Kale HW II and Maehr DS. 1990. Florida's Birds: A Handbook and Reference. Sarasota: Pineapple Press. 288 p.
- Kushlan, J. A. 1990. Freshwater marshes. Pages 324-363 in R. L. Myers and J. J. Ewel editors. Ecosystems of Florida. University of Central Florida Press; Orlando, Florida.
- Langeland, K. A., and R. K. Stocker. 2001. Control of non-native plants in natural areas of Florida. University of Florida Cooperative Extension Service Document SP 242. 34pp. University of Florida, UF/IFAS Extension Digital Information Source (EDIS)

 Database.


 Available from http://edis.ifas.ufl.edu/pdffiles/WG/WG20900.pdf (accessed January 2010).
- Langeland, K. A., D. W. Clark, R. K. Stocker, and G. E. MacDonald. 2003. Evaluation of Foliar Applied Herbicides for Control of Christmas Senna. In Wildland Weeds. Winter 2009. Available from http://www.se-eppc.org/wildlandweeds/pdf/Winter2003-Langeland-pp13-14.pdf (accessed January 2010).

102

- Langeland, K. A. 2008. Natural area weeds: distinguishing native and non-native "Boston ferns" and "sword ferns" (*Nephrolepis* spp.). University of Florida Cooperative Extension Service Document SS-AGR-22. 7pp. University of Florida, UF/IFAS Extension Digital Information Source (EDIS) Database. Available from http://edis.ifas.ufl.edu/pdffiles/AG/AG12000.pdf (accessed January 2010).
- Larson, B. C., J. H. Frank, G. M. Allen, M. B. Main. 2006. Florida's native bromeliads. University of Florida Cooperative Extension Service Circular 1466. 10pp. University of Florida, UF/IFAS Extension Digital Information Source (EDIS) Database. Available from http://edis.ifas.ufl.edu/UW205 (accessed November 2007).
- Lee, J. C. 1985. *Anolis sagrei* in Florida: phenetics of a colonizing species I. Meristic characters. Copeia 1985:182-194.
- Lodge, T. E. 2005. The Everglades handbook Understanding the Ecosystem. 2nd edition. CRC Press, Boca Raton, FL.
- Liudahl, K., D.J. Belz, L. Carey, R.W. Drew, S. Fisher, and R. Pate. 1990. Soil Survey of Collier County Area Florida. USDA, Natural Resources Conservation Service; Washington, D.C.
- Mazziotti, F. J. 2002. Wood Storks (Mycteria Americana). Wildlife Ecology and Conservation Department, Florida Cooperative Extension Service document SSWIS12. 2pp. University of Florida, UF/IFAS Extension Digital Information Source (EDIS) Database. Available from http://edis.ifas.ufl.edu/UW065 (accessed December 2009).
- Miller J. A. 1986. Hydrogeologic framework of the Floridan Aquifer System in Florida and in parts of Georgia, Alabama, and South Carolina. United States Geological Survey Professional Paper 1403-B. United States Government Printing Office, Washington, D.C.
- Natural Resource Conservation Service (NRCS). 2003 Feb. Conservation Practice Standard Prescribed Burning. Field Office Technical Guide Section IV. (FL): Code 338.7 p.
- Nelson, G. 2000. The ferns of Florida: a reference and field guide. Pineapple Press, Sarasota, FL.
- Oaks, R. Q. and J. R. Dunbar. 1974. Post Miocene stratigraphy of the Central and Southern Atlantic Coastal Plain. Utah State University Press, Logan, Utah.

- Schwartz, A. and R. W. Henderson. 1991. Amphibians and reptiles of the West Indies: descriptions, distributions, and natural history. University of Florida Press, Gainesville.
- Scott, T. M. 1988. Lithostratigraphy of the Hawthorne Group (Miocene). Florida Geological Survey Bulletin No. 59, Tallahassee, Florida.
- Stokes, C. 2009. From Crop to Weed Natalgrass in Florida. Pp. 8-9. In Wildland Weeds. Summer 2009, Volume 12, Number 3. Florida Exotic Pest Plant Council. Gainesville, FL.
- Thornton, D. H., M.E. Sunquist, M.B. Main 2004. Ecological Separation within Newly Sympatric Populations of Coyotes and Bobcats in South Central Florida. Journal of Mammalogy 85(5) 973-982.
- United States Department of Agriculture (USDA)/Soil Conservation Service. 1984. Soil Survey of Collier County.
- United States Department of the Interior (USDI). 2004. U.S. Geological Services historic aerial photo web page available from http://sofia.usgs.gov/publications/ofr/02-327/htm/imagedir.htm (accessed on December 2009).
- United States Fish and Wildlife Service (USFWS). 1999. Mesic temperate hammocks. South Florida multi-species recovery plan a species plan…an ecosystem approach. USFWS Southeast Region, Compact Disk.
- United States Fish and Wildlife Service (USFWS). 1999. Mesic flatwoods. South Florida multi-species recovery plan a species plan…an ecosystem approach. USFWS Southeast Region, Compact Disk.
- United States Fish and Wildlife Service (USFWS). 2009. Federally listed species list available from http://www.fws.gov/endangered/wildlife.html#Species (accessed on December 2009).
- Wade D. D., and J. D. Lundsford. 1989. A guide for prescribed fire in southern forests. National Wildfire Coordinating Group. Technical Publication R8-TP 11. 56 p. Available from: National Interagency Fire Center, ATTN: Supply, 3833 S. Development Ave., Boise, ID 83705. Order NFES #2108.
- Wunderlin, R. P., and B. F. Hansen. 2008. Atlas of Florida vascular plants. [S.M. Landry and K.N. Campbell (application development), Floirda Center for Community Design and Research]. Institute for Systematic Botany, University of South Florida, Tampa. Available from http://www.plantatlas.usf.edu/.

Pepper Ranch Preserve Legal Description





Appendix 2 Florida Land Use, Cover and Forms Classification System Designations for Pepper Ranch Preserve – (2'x3' map)


(Same as map on page 22 only larger)

Florida Natural Areas Inventory Designations for Pepper Ranch Preserve – (2'x3' map)

(Same map as on Page 24 only larger)







Cattle Lease

Pepper Ranch Preserve Quality Wildlife Management Hunt Program 2010-2011 Brochure

Pepper Ranch Preserve – Land Use Compatibility Matrix

	Pepper Ranch Pres	erve - (Compat	ibility N	latrix														
	PUBLIC USES										MITIGATION & LAND USE				OTHER REVENUE GENERATING		Conservation		
		hiking	mountain biking	primitive camping	horseback riding	hunting	fishing	nature photography	special events lodge rental	ecotourism	campground	archery	Panther Conservatio n Bank	wetlands mitigation	water storage	SSA	oil drilling	cattle grazing	Collier Ordinance No. 2007-65
PUBLIC USES	hiking																		
	mountain biking																		
	primitive camping																		
	horseback riding																		
	hunting																		
	fishing																		
	nature photography																		
	special events logde rental																		
	ecotourism																		
	campground																		
	archery																		
MITIGATION & LAND USE	Panther Conservation Bank																		
	Wetlands Mitigation																		
	water storage																		
	SSA																		
OTHER REVENUE GENERATING	oil drilling																		
	cattle grazing																		
Conservat	ion Collier Ordinance No. 2007- 65																		
		Uses are compatible throughout the Preserve Uses are compatible but on seperate portions of the Preserve																	
					ible during c														

Appendix 9	
Parcel Folio Map of Pepper Ranch Preserve	

